



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|---|
| (51) International Patent Classification ⁶ : C12N 9/78, 15/55, 15/31, C07K 14/21, C12N 15/63, 15/62, 1/19, 1/21, C12P 13/02, 7/40 // (C12N 9/78, C12R 1:40) (C12N 1/19, C12R 1:84) | A2 | (11) International Publication Number: WO 97/12964 (43) International Publication Date: 10 April 1997 (10.04.97) |
| (21) International Application Number: PCT/US96/15969 (22) International Filing Date: 3 October 1996 (03.10.96) (30) Priority Data: 60/004,914 6 October 1995 (06.10.95) US (71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FALLON, Robert, Donald [US/US]; 104 Black Oak Drive, Elkton, MD 21921- 2022 (US). NELSON, Mark, James [US/US]; 707 Fiske Lane, Newark, DE 19711-3133 (US). PAYNE, Mark, Scott [US/US]; 2408 Newell Drive, Wilmington, DE 19808-3344 (US). (74) Agents: FLOYD, Linda, A. et al.; E.I. du Pont de Nemours and Company, Legal Patent Record Center, 1007 Market Street, Wilmington, DE 19898 (US). | | (81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i> |
| (54) Title: NUCLEIC ACID FRAGMENTS ENCODING STEREOSPECIFIC NITRILE HYDRATASE AND AMIDASE ENZYMES AND RECOMBINANT MICROORGANISMS EXPRESSING THOSE ENZYMES USEFUL FOR THE PRODUCTION OF CHIRAL AMIDES AND ACIDES (57) Abstract The present invention provides a nitrile hydratase nucleic acid fragment isolated from <i>Pseudomonas putida</i> which encodes a nitrile hydratase activity capable of catalyzing the hydrolysis of certain racemic nitriles to the corresponding R- or S- amides. Also provided are transformed microorganisms capable of the active expression of said nitrile hydratase activity. Additionally, the invention provides a transformant harboring the nitrile hydratase gene in conjunction with an amidase gene, both of which may be co-expressed producing active nitrile hydratase and amidase enzymes respectively. Methods for the production of such enantiomeric materials are also provided. | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AM | Armenia | GB | United Kingdom | MW | Malawi |
| AT | Austria | GE | Georgia | MX | Mexico |
| AU | Australia | GN | Guinea | NE | Niger |
| BB | Barbados | GR | Greece | NL | Netherlands |
| BE | Belgium | HU | Hungary | NO | Norway |
| BF | Burkina Faso | IE | Ireland | NZ | New Zealand |
| BG | Bulgaria | IT | Italy | PL | Poland |
| BJ | Benin | JP | Japan | PT | Portugal |
| BR | Brazil | KE | Kenya | RO | Romania |
| BY | Belarus | KG | Kyrgyzstan | RU | Russian Federation |
| CA | Canada | KP | Democratic People's Republic of Korea | SD | Sudan |
| CF | Central African Republic | KR | Republic of Korea | SE | Sweden |
| CG | Congo | KZ | Kazakhstan | SG | Singapore |
| CH | Switzerland | LJ | Liechtenstein | SI | Slovenia |
| CI | Côte d'Ivoire | LK | Sri Lanka | SK | Slovakia |
| CM | Cameroon | LR | Liberia | SN | Senegal |
| CN | China | LT | Lithuania | SZ | Swaziland |
| CS | Czechoslovakia | LU | Luxembourg | TD | Chad |
| CZ | Czech Republic | LV | Larvia | TG | Togo |
| DE | Germany | MC | Monaco | TJ | Tajikistan |
| DK | Denmark | MD | Republic of Moldova | TT | Trinidad and Tobago |
| EE | Estonia | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | UG | Uganda |
| FI | Finland | MN | Mongolia | US | United States of America |
| FR | France | MR | Mauritania | UZ | Uzbekistan |
| GA | Gabon | | | VN | Viet Nam |

TITLE

NUCLEIC ACID FRAGMENTS ENCODING STEREOSPECIFIC NITRILE HYDRATASE AND AMIDASE ENZYMES AND RECOMBINANT MICROORGANISMS EXPRESSING THOSE ENZYMES USEFUL FOR THE PRODUCTION OF CHIRAL AMIDES AND ACIDS


FIELD OF INVENTION

The present invention relates to the field of molecular biology and methods for the isolation and expression of foreign genes in recombinant microorganisms. More specifically, the invention relates to the isolation, sequencing, and recombinant expression of nucleic acid fragments (genes) encoding a stereospecific, nitrile hydratase (NHase) activity capable of catalyzing the hydrolysis of certain racemic nitriles to the corresponding R- or S- amides. Additionally, the invention relates to the co-expression of the nitrile hydratase nucleic acid fragment with a nucleic acid fragment encoding a stereospecific amidase activity capable of converting a racemic mixture of R- and S- amides to the corresponding enantiomeric R- or S- carboxylic acids.


BACKGROUND

Many agrochemicals and pharmaceuticals of the general formula X-CHR-COOH are currently marketed as racemic or diastereomer mixtures. In many cases the physiological effect derives from only one enantiomer/diastereomer where the other enantiomer/diastereomer is inactive or even harmful. Methods for synthesizing enantiomers are becoming increasingly important tools for the production of chemicals of enantiomer purity. To date, however, no recombinant, stereospecific NHase has been described capable of catalyzing the hydrolysis of certain racemic nitriles to the corresponding R- or S- amides. Methods for the selective preparation of stereospecific amides from nitriles are known and incorporate microorganisms possessing nitrile hydratase activity (NHase). These NHases catalyze the

addition of one molecule of water to the nitrile, resulting in the formation of the amide free product according to Reaction 1:

Reaction 1

Similarly, methods for the stereospecific
5 production of carboxylic acids are known and incorporate microorganisms possessing an amidase (Am) activity. In general amidases convert the amide product of Reaction 1 to the acid free product plus ammonia according to Reaction 2:

Reaction 2

10 A wide variety of bacterial genera are known to possess a diverse spectrum of nitrile hydratase and amidase activities including *Rhodococcus*, *Pseudomonas*, *Alcaligenes*, *Arthrobacter*, *Bacillus*, *Bacteridium*, *Brevibacterium*, *Corynebacterium*, and *Micrococcus*. For
15 example, nitrile hydratase enzymes have been isolated from *Pseudomonas chlororaphis*, B23 [Nishiyama, M. J., *Bacteriol.*, 173:2465-2472 (1991)] *Rhodococcus rhodochromis* J1 [Kobayashi, M., *Biochem. Biophys. Acta*, 1129:23-33 (1991)] *Brevibacterium sp.* 312 (Mayaux
20 et al., *J. Bacteriol.*, 172:6764-6773 (1990)), and *Rhodococcus sp.* N-774 [Ikehata, O., Nishiyama, M., Horinouchi, S., Beppu, T., *Eur. J. Biochem.*, 181: 563-570 (1989)]. No disclosure of any stereoselective activity is made for any of these enzymes. Only two
25 disclosures have been made for stereoselective nitrile hydratase activity in native bacterial strains. The Applicants have disclosed a stereospecific nitrile hydratase from *P. putida* NRRL-18668 [WO 92/05275 (1990)].

30 Wildtype microorganisms known to possess nitrile hydratase activity have been used to convert nitriles to amides and carboxylic acids. For example, EPA 326,482 discloses the stereospecific preparation of aryl-2-alkanoic acids such as 2-(4-chlorophenyl)-3-

methybutyric acid by microbial hydrolysis of the corresponding racemic amide using members of *Brevibacterium* and *Corynebacterium*. Similarly, U.S. Pat. No. 4,366,250 teaches the use of *Bacillus*,
5 *Bacteridium*, *Micrococcus* and *Brevibacterium* in a method for the preparation of L-amino acids from the corresponding racemic amino nitriles. Finally, WO 92/05275 teaches a biologically-catalyzed method for converting a racemic alkyl nitrile to the
10 corresponding R- or S-alkanoic acid through an intermediate amide using members of the bacterial genera *Pseudomonas* spp. (e.g., *putida*, *aureofaciens*, *Moraxella* spp.) and *Serratia* (e.g., *Serratia liquefaciens*).
15 In addition to the use of wildtype organisms, recombinant organisms containing heterologous genes for the expression of nitrile hydratase are also known for the conversion of nitriles. For example, Cerebelaud et al., (WO 9504828) teach the isolation
20 and expression in *E. coli* of nitrile hydratase genes isolated from *C. testosteroni*. The transformed hosts effectively convert nitriles to amides where the nitrile substrate consists of one nitrile and one carboxylate group. However, WO 9504828 does not teach
25 a stereospecific conversion of nitriles.

Similarly, Beppu et al., (EP 5024576) disclose plasmids carrying both nitrile hydratase and amidase genes from *Rhodococcus* capable of transforming *E. coli* where the transformed host is then able to use
30 isobutyronitrile and isobutyroamide as enzymatic substrates. However, EP 5024576 does not teach a stereospecific conversion of nitriles or amides.

As with nitrile hydratases, microorganisms possessing amidase activity have been used to convert
35 amides to carboxylic acids. In USSN 08/403911, Applicants disclose a method for converting an (S)-amide, or stereospecifically converting a mixture of (R)- and (S)-amides to the corresponding

enantiomeric (S)-carboxylic acid by contacting said amide with *Pseudomonas chlororaphis* B23 in a solvent. This method uses a wildtype microorganism and does not anticipate a recombinant catalyst or heterologous gene expression. Blakey et al., *FEMS Microbiology Letters*, 129:57-62 (1995) disclose a *Rhodococcus* sp. having activity against a broad range of nitriles and dinitriles and able to catalyze regio-specific and stereo-specific nitrile biotransformations.

Genes encoding amidase activity have been cloned, sequenced, and expressed in recombinant organisms. For example, Azza et al., (*FEMS Microbiol. Lett.* 122, 129, (1994)) disclose the cloning and over-expression in *E. coli* of an amidase gene from *Brevibacterium* sp. R312 under the control of the native promoter. Similarly, Kobayashi et al., (*Eur. J. Biochem.*, 217, 327, (1993)) teach the cloning of both a nitrile hydratase and amidase gene from *R. rhodococcus* J1 and their co-expression in *E. coli*.

What is needed and inventive over the prior art is a method for the stereospecific conversion of racemic alkyl nitriles to the corresponding R- or S-alkanoic acids using a recombinant organism.

SUMMARY OF THE INVENTION

This invention relates to nucleic acid fragments encoding:

- 1) the α subunit of a stereospecific nitrile hydratase enzyme, said gene having at least a 64% base homology with the α subunit coding region of the *Rhodococcus rhodochrous* J1 L-NHase gene [Kobayashi, M., *Biochem. Biophys. Acta*, 1129:23-33 (1991)] and said enzyme capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides; and
- 2) the β subunit of a stereospecific nitrile hydratase enzyme, said gene having at least a 52% base homology with the β subunit coding region of the *Rhodococcus rhodochrous* J1 L-NHase gene and said

enzyme capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S-amides.

Another embodiment of the invention is a nucleic acid fragment comprising the nucleic acid fragments encoding both the α and β subunits of a stereospecific nitrile hydratase enzyme described above, said enzyme capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S-amides.

A further embodiment of the invention is a nucleic acid fragment encoding the α subunit of a stereospecific nitrile hydratase enzyme, said nucleic acid fragment having the nucleotide sequence as represented in SEQ ID NO.:3 and said enzyme capable of catalyzing the hydrolysis of racemic alkyl nitriles to the corresponding R- or S- amides.

A further embodiment of the invention is a nucleic acid fragment encoding the β subunit of a stereospecific nitrile hydratase enzyme, said nucleic acid fragment having the nucleotide sequence as represented in SEQ ID NO.:4 and said enzyme capable of catalyzing the hydrolysis of racemic alkyl nitriles to the corresponding R- or S- amides.

Still another embodiment of the invention is a nucleic acid fragment encoding both the α and β subunits of a stereospecific nitrile hydratase enzyme, said nucleic acid fragment having the nucleotide sequence as represented in SEQ ID NO.:17 and said enzyme capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.

Further embodiments of the invention include

1) the polypeptide α subunit of a stereospecific nitrile hydratase enzyme, said α subunit having the amino acid sequence as represented in SEQ ID NO.:1 and said enzyme being capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides; and

2) the polypeptide β subunit of a stereospecific nitrile hydratase enzyme, said β subunit having the amino acid sequence as represented in SEQ ID NO.:2 and said enzyme being capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.

A further embodiment of the invention is a stereospecific nitrile hydratase enzyme, said enzyme comprising the combined α and β subunits having the respective amino acid sequences SEQ ID NOs.:1 and 2 in proper conformation such that said enzyme catalyzes the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.

A still further embodiment of the invention is a 6.5 kb nucleic acid fragment encoding a nitrile hydratase enzyme and the accessory nucleic acid fragments necessary for the enzymes's active expression and further characterized by the restriction fragment map shown in Figure 2. This 6.5 kb nucleic acid fragment is incorporated into an expression vector capable of transforming a suitable host cell for the expression of active stereospecific nitrile hydratase as characterized by the plasmid map shown in Figure 3.

The invention further provides a region of the *P. putidia* genome encompassed within the 6.5 kb fragment, designated P14K, which encodes a polypeptide that is necessary for the bioactivity of the stereospecific nitrile hydratase enzyme isolated from *Pseudomonas putida* NRRL-18668.

Additionally the invention provides a nucleic acid fragment encoding a 18668 amidase having an amino acid sequence as represented in SEQ ID NO.:28, wherein the amino acid sequence may encompass amino acid substitutions, deletions or additions that do not alter the function of said amidase. The 18668 amidase is isolated from *Pseudomonas putida* NRRL-18668 and is

distinct from the amidase isolated from *Pseudomonas chlororaphis* B-23 (FERM B-187).

The present invention further provides recombinant hosts, transformed with the nucleic acid
5 fragment encoding a 18668 amidase and/or the genes encoding the α , β nitrile hydratase subunits and the P14K region of the *Pseudomonas putida* NRRL-18668 genome.

The invention also provides methods for the
10 conversion of racemic nitriles to the corresponding R- or S-amides or corresponding enantiomeric R- or S-carboxylic acids using the above transformed hosts containing nucleic acid fragments encoding a 18668 amidase and/or the genes encoding the α , β nitrile
15 hydratase subunits and the P14K region of the *Pseudomonas putida* NRRL-18668 genome.

Other embodiments of the invention are:

- 1) a transformed microbial host cell comprising the nucleic acid fragment represented by
20 SEQ ID NO.:17 wherein said host cell expresses active nitrile hydratase enzyme capable of catalyzing the hydrolysis of racemic aryl-2 alkane nitriles to the corresponding R- or S- amides; and
- 2) a transformed microbial host cell
25 comprising the 6.5 kb nucleic acid fragment characterized by the restriction map shown in Figure 2 wherein said host cell expresses active nitrile hydratase enzyme capable of catalyzing the hydrolysis of racemic aryl-2 alkane nitriles to the corresponding
30 R- or S- amides.


Other embodiments of the invention are host cells transformed with nucleic acid fragments represented by SEQ ID NO.:17 or the restriction maps of Figures 2 and 3, wherein the host cell is selected from the group
35 consisting of bacteria of the genera *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, and *Streptomyces*, yeast of the genera *Pichia*, *Hansenula*,

and *Saccharomyces*, and filamentous fungi of the genera *Aspergillus*, *Neurospora*, and *Penicillium*.

A particular embodiment of the invention is *Escherichia coli* transformed with the nucleic acid
 5 fragment represented by SEQ ID NO.:17 or the nucleic acid fragment represented by the restriction map of Figure 2.






A further embodiment of the invention is an expression vector described in Figure 6 comprising
 10 1) a 5.0 kb nucleic acid fragment from the 6.5 kb fragment of Claim 10, and 2) a nucleic acid fragment having the nucleic acid sequence as given in SEQ ID NO.:20, wherein said nucleic acid fragment encodes an amidase enzyme, and wherein said expression vector is
 15 capable of transforming suitable host cells for the co-expression of active stereospecific nitrile hydratase and amidase. A further embodiment is a host cell transformed with this expression vector wherein more particularly the host is selected from the group
 20 consisting of the genera *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Streptomyces*, *Hansenula*, *Saccharomyces*, *Pichia*, *Aspergillus*, *Neurospora*, and *Penicillium*. A further embodiment is *Escherichia coli* SW17 transformed with pSW17.

25 A further embodiment of the invention is a method for converting a nitrile of the formula



wherein:


A is selected from the group consisting of:




R¹ is C₁-C₄ alkyl;

R² is H or OH;

R³ is H, Cl, OCF₂H, (CH₃)₂CHCH₂, H₂C=C(CH₃)CH₂NH,





R^4 is Cl or F;

to the corresponding amide comprising contacting said nitrile with the transformed host cell containing a nucleic acid fragment having the nucleotide sequence represented by SEQ ID NO.:17 that stereospecifically converts the racemic nitrile to the corresponding enantiomeric R- or S-amide, the host cell selected from the group consisting of *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Streptomyces*, *Hansenula*, *Saccharomyces*, *Pichia*, *Aspergillus*, *Neurospora*, and *Penicillium*.

The Applicants also provide a method for the conversion of the above described nitrile to corresponding enantiomeric (R) or (S)-carboxylic acid by contacting the nitrile with the transformed host comprising an expression vector comprising a nucleic acid fragment represented by Figure 2 and the nucleic acid sequence of SEQ ID NO.:20, the host cell selected from the group consisting of *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Streptomyces*, *Hansenula*, *Saccharomyces*, *Pichia*, *Aspergillus*, *Neurospora*, and *Penicillium*.

A further embodiment of the invention is a nucleic acid fragment encoding the α and β subunits of a stereospecific nitrile hydratase enzyme, said portion of the nucleic acid fragment encoding the α subunit having at least a 64% base homology to the *Rhodochrous* J1 L-NHase gene and said portion of the nucleic acid fragment encoding the β subunit having a 52% base homology to the *Rhodochrous* J1 L-NHase gene, and said enzyme capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.

Yet another embodiment of the invention is the polypeptide encoded by any one of the nucleic acid fragments of the invention.

Embodiments of the invention are plasmids pSW2
5 carried in SW2 and designated as ATCC 69888, pSW17
carried in SW17 and designated as ATCC 69887, pSW50
carried in *P. pastoris* SW50.2 and designated as
ATCC 74391, pSW37 carried in *E. coli* SW37 and
designated as ATCC 98174, and pSW23 carried in *E. coli*
10 SW23 and designated as ATCC 98175.

BRIEF DESCRIPTION OF THE FIGURES

BIOLOGICAL DEPOSITS AND SEQUENCE LISTING

Figure 1 is a plasmid map of the plasmid pSW1
containing a 6.5 kb DNA fragment which encodes the α
15 and β subunits of the nitrile hydratase enzyme
isolated from *P. putida* (NRRL-18668).

Figure 2 is a restriction map of the 6.5 kb
nucleic acid fragment which includes the nitrile
hydratase gene isolated from *P. putida* (NRRL-18668)
20 showing the location of the α and β subunits.

Figure 3 is a plasmid map of the plasmid pSW2
created by inserting the 6.5 kb DNA fragment
comprising the genes encoding the α and β subunits of
nitrile hydratase into the wide-host-range vector
25 pMMB207.

Figure 4 is a plasmid map of the plasmid pSW5
created by inserting a 2.8 kb subclone of the 6.5 kb
nucleic acid fragment comprising the genes encoding
the α and β subunits of nitrile hydratase into the wide-
30 host-range vector pMMB207.

Figure 5 is a western blot analysis showing the
production of NRRL-18668 nitrile hydratase protein in
E. coli. (A) Coomassie Blue stained SDS-PAGE gel of
protein extracts from uninduced (u) and induced (i)
35 *E. coli* transformed with the plasmid pSW2.
(B) Western blot analysis of duplicate gel shown in
(A) using anti-NH sera. M, protein molecular weight

markers; NH, nitrile hydratase protein from NRRL-18668. Arrow indicates NH.

Figure 6 is a plasmid map of the plasmid pSW17 created by inserting a 1.5 kb DNA fragment comprising the gene encoding amidase from *Pseudomonas chlororaphis* B23, and a 5.0 kb subclone of the 6.5 kb DNA fragment comprising the genes encoding the α and β subunits of nitrile hydratase into the wide-host-range vector pMMB207.

Figure 7 illustrates the nucleotide and amino acid sequences of the *Pseudomonas putida* (NRRL-18668) α and β nitrile hydratase coding regions also found in SEQ ID NO.:17.

Figure 8 is a restriction map of the 6.5 kb nucleic acid fragment which includes the nitrile hydratase gene isolated from *P. putida* (NRRL-18668) plus sequence upstream of the EcoR1 site (shown in Figure 2) including a new Pst1 site.

Figure 9 is a restriction map of the 6.5 kb nucleic acid fragment which includes the nitrile hydratase gene isolated from *P. putida* (NRRL-18668) plus sequence upstream of the new Pst1 site (shown in Figure 8) including a new EcoR1 site.

Figure 10 is a restriction map of an 8 kb nucleic acid fragment showing the 6.5 kb nucleic acid fragment which includes the nitrile hydratase gene isolated from *P. putida* (NRRL-18668), P14K, and the region encoding a *P. putida* (NRRL-18668) amidase enzyme.

Figure 11 is a plasmid map of pHIL-D4B2 created by replacing the 0.9 kb EcoR1/Xba1 fragment in pHIL-D4 with the 0.9 kb EcoR1/Xba1 fragment from pAO815.

Figure 12 is a plasmid map of pSW46 created by the insertion of the α gene of the nitrile hydratase enzyme into the EcoR1 site of pHIL-D4B2.

Figure 13 is a plasmid map of pSW47 created by the insertion of the β gene of the nitrile hydratase enzyme into the EcoR1 site of pHIL-D4B2.

Figure 14 is a plasmid map of pSW48 created by the insertion of the P14K gene into the EcoR1 site of PHIL-D4B2.

Figure 15 is a plasmid map of pSW49 containing the α and β expression cassettes from pSW46 and pSW47.

Figure 16 is a plasmid map of pSW50 containing the α , β and P14K expression cassettes from pSW46, pSW47 and pSW48.

Figure 17 is a plasmid map of pSW37 containing the expression cassette for the amidase isolated from *P. putida* (NRRL-18668).

Figure 18 is a plasmid map of pSW23 containing the expression cassette for the amidase, α , β and P14K isolated from *P. putida* (NRRL-18668).

Applicants have provided sequence listings 1-28 in conformity with 37 C.F.R. 1.821-1.825 and Appendices A and B ("Requirements for Application Disclosures Containing Nucleotides and/or Amino Acid Sequences") and in conformity with "Rules for the Standard Representation of Nucleotide and Amino Acid Sequences in Patent Applications" and Annexes I and II to the Decision of the President of the EPO, published in Supplement No. 2 to OJ EPO, 12/1992.

Applicants have made the following biological deposits under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for the Purposes of Patent Procedure:

| Depositor Identification Reference | Int'l. Depository Designation | Date of Deposit |
|--|-------------------------------|-------------------|
| <i>Pseudomonas Putida</i> | NRRL 18668 | 6 July 1990 |
| <i>Escherichia coli</i> SW2 carrying pSW2 | ATCC 69888 | 15 August 1995 |
| <i>Escherichia coli</i> SW17 carrying pSW17 | ATCC 69887 | 15 August 1995 |
| <i>Pichia pastoris</i> SW50.2 carrying pSW50 | ATCC 74391 | 20 September 1996 |
| <i>E. coli</i> SW37 carrying pSW37 | ATCC 98174 | 20 September 1996 |
| <i>E. coli</i> SW23 carrying pSW23 | ATCC 98175 | 20 September 1996 |

As used herein, "NRRL" refers to the Northern Regional Research Laboratory, Agricultural Research Service Culture Collection International Depository

Authority located at 11815 N. University Street,
Peoria, IL 61604 U.S.A. The "NRRL No." is the
accession number to cultures on deposit at the NRRL.

As used herein, "ATCC" refers to the American
5 Type Culture Collection International Depository
Authority located at 12301 Parklawn Drive, Rockville,
MD 20852 U.S.A. The "ATCC No." is the accession
number to cultures on deposit with the ATCC.

DETAILED DESCRIPTION OF THE INVENTION

10 The present invention provides genes derived from
Pseudomonas putida (NRRL-18668) which encode two
polypeptides, which, in combination, have the ability
to act as a catalyst to selectively hydrate one
nitrile enantiomer in a racemic mixture to produce the
15 chiral amide. This invention also provides a
recombinant nucleic acid fragment containing the genes
and a set of transformed microbial cell hosts
containing the recombinant nucleic acid fragment. The
invention further provides a method for the production
20 of the polypeptide catalysts using the transformed
microbes and the use of the catalyst in chiral amide
production. Additionally, the invention provides for
the co-expression in a transformed host of the nitrile
hydratase genes with the genes encoding a
25 stereospecific amidase derived from *Pseudomonas*
chlororaphis B-23 (FERM B-187) for the production of
chiral acids.

The following definitions are used herein and
should be referred to for interpretation of the claims
30 and the specification.


Abbreviations:

| | | |
|----|-------|--|
| | CPIA | - 2-(4-chlorophenyl)-3-methylbutyric acid |
| | CPIAm | - 2-(4-chlorophenyl)-3-methylbutyramide |
| | CPIN | - 2-(4-chlorophenyl)-3-methylbutyronitrile |
| 35 | GC | - Gas Chromatography |
| | HPLC | - High-Performance Liquid Chromatography |
| | IPTG | - isopropyl-b-D-thiogalactopyranoside |

SDS Page - Sodium dodecyl sulfate polyacrylimide
gel electrophoresis


The term "nitrile hydratase" refers to an enzyme
isolated from the bacteria *Pseudomonas putida*


- 5 (NRRL-18668) which is characterized by its ability to
convert a racemic alkyl nitrile to the corresponding
enantiomeric R- or S-amide through an intermediate
amide where the starting nitrile is:




and wherein:

- 10 A is selected from the group consisting of:






A-8




A-9



A-10

and








A-11


R^1 is C_1 - C_4 alkyl;

R^2 is H or OH;

R^3 is H, Cl, OCF_2H , $(CH_3)_2CHCH_2$, $H_2C=C(CH_3)CH_2NH$,



or




; and

R^4 is Cl or F.

5 More specifically, the enzyme has an ability to connect the racemic alkyl nitrile to the corresponding enantiomeric R- or S-alkanoic acid through an intermediate amide.

10 The instant nitrile hydratase is further defined by the amino acid sequences of its α and β subunits as respectively given in SEQ ID NO.:1 and SEQ ID NO.:2 which are encoded by the α and β nitrile hydratase subunit genes whose base sequences are respectively given by SEQ ID NO.:3 and SEQ ID NO.:4.


15 The term "amidase" refers to an enzyme naturally found in the bacterium *Pseudomonas putida* B23 (FERM B-187) which is characterized by its ability to convert amides of the structure:



I


wherein:

A is selected from the group consisting of:



A-1

and



A-2

R^1 is C_1 - C_4 alkyl;

R^2 is H; F; Cl; Br; OH; C_1 - C_3 alkyl; OCF_2H ; or
 $\text{H}_2\text{C}=\text{C}(\text{CH}_3)\text{CH}_2\text{NH}$; and

R^3 is H; F; Cl; Br; OH; C_1 - C_3 alkyl; or C_1 - C_3
 alkoxy;

to the corresponding enantiomeric (R) or
 (S)-carboxylic acid. The amidase of the instant
 invention is further identified by the amino acid
 sequence given in Nishiyama et al., *Bacterial.*,
 173:2465-2472 (1991) and the DNA base sequence
 disclosed in SEQ ID NO.:20.

The term "18668 amidase" refers to an enzyme
 naturally found in the bacterium *Pseudomonas putida*
 NRRL-18668 which is characterized by its ability to
 convert C3 to C6 amides to the corresponding acids.
 In addition, as described in PCT/DK91/00189, the 18668
 amidase is characterized by the ability to convert
 some (R,S)-aryl-2-alkane nitriles to the corresponding
 enantiomerically enriched (R) or (S)-carboxylic acid.
 The amidase of the instant invention is further
 identified by the amino acid sequence given in SEQ ID
 NO.:28 and the DNA base sequence disclosed in SEQ ID
 NO.:27. The "18668 amidase" is distinct from the
 amidase isolated from bacterium *Pseudomonas putida*
 B23 (FERM B-187).

The term "P14K gene" refers to a region of the *Pseudomonas putida* NRRL-18668 genome encoding a polypeptide as given by SEQ ID NO.:22 having the base sequence as given by SEQ ID NO.:21, where the
5 expression of the P14K gene is essential for the bioactivity of the *Pseudomonas putida* NRRL-18668 nitrile hydratase enzyme. The term "P14K polypeptide" (or "P14K protein") refers to the active polypeptide encoded by the P14K region.

10 "Transformation" refers to the acquisition of new genes in a cell by the incorporation of nucleic acid.

The term "nucleic acid" refers to complex compounds of high molecular weight occurring in living cells, the fundamental units of which are nucleotides
15 linked together with phosphate bridges. Nucleic acids are subdivided into two types: ribonucleic acid (RNA) and deoxyribonucleic acid (DNA).

The terms "host cell" and "host organism" refer to a microorganism capable of incorporating foreign or
20 heterologous genes and expressing those genes to produce an active gene product.

The terms "foreign gene", "foreign DNA", "heterologous gene", and "heterologous DNA" refer to genetic material native to one organism that has been
25 placed within a host organism.

The terms "recombinant organism", "transformed host", and "transformed microbial host" refer to an organism having been transformed with heterologous or
30 foreign genes. The recombinant organisms of the present invention express foreign genes encoding active nitrile hydratase and amidase enzymes.

The term "nucleic acid fragment" refers to a fragment of DNA that may encode a gene and/or regulatory sequences preceding (5' non-coding) and
35 following (3' non-coding) the coding region (gene).

The term "expression" refers to the transcription and translation to gene product from a gene coding for the sequence of the gene product, usually a protein.

The terms "plasmid" and "vector" refer to an extra chromosomal element often carrying genes which are not part of the central metabolism of the cell, and usually in the form of circular double-stranded DNA molecules. Such elements may be autonomously replicating sequences, genome integrating sequences, phage sequences, linear or circular, of a single- or double-stranded DNA or RNA, derived from any source.

The term "cassette" refers to a number of nucleotide sequences which have been joined or recombined into a unique construction. An "expression cassette" is specifically comprised of a promoter fragment, a DNA sequence for a selected gene product, and a transcriptional termination sequence.

The terms "restriction endonuclease" and "restriction enzyme" refer to an enzyme which catalyzes hydrolytic cleavage within a specific nucleotide sequence in double-stranded DNA.

The term "promoter" refers to a sequence of DNA, usually upstream of (5' to) the protein coding sequence of a structural gene, which controls the expression of the coding region by providing the recognition for RNA polymerase and/or other factors required for transcription to start at the correct site.

A "fragment" constitutes a fraction of the complete nucleic acid sequence of a particular region. A fragment may constitute an entire gene.

The terms "peptide", "polypeptide" and "protein" are used interchangeably to refer to the gene product expressed.

The terms "encoding" and "coding" refer to the process by which a gene, through the mechanisms of transcription and translation, produces an amino acid sequence. The process of encoding a specific amino acid sequence includes DNA sequences that may involve base changes that do not cause a change in the encoded amino acid, or which involve base changes which may

alter one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence. It is therefore understood that the invention encompasses more than the specific exemplary sequences. Modifications to the sequence, such as deletions, insertions, or substitutions in the sequence which produce silent changes that do not substantially affect the functional properties of the resulting protein molecule are also contemplated. For example, alteration in the gene sequence which reflect the degeneracy of the genetic code, or which result in the production of a chemically equivalent amino acid at a given site, are contemplated. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a biologically equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the protein molecule would also not be expected to alter the activity of the protein. In some cases, it may, in fact, be desirable to make mutants of the sequence in order to study the effect of alteration on the biological activity of the protein. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity in the encoded products. Moreover, the skilled artisan recognizes that sequences encompassed by this invention are also defined by their ability to hybridize, under stringent conditions (0.1X SSC, 0.1% SDS, 65°C), with the sequences exemplified herein.

"Homology" refers to the degree to which two nucleic acid fragments contain the same base sequence. "Homology" is determined by the operation of an algorithm and is expressed as a percentage of the
5 base sequence that is the same in both fragments.

Applicants have accomplished the following which are discussed in more detail below and in the Examples:

I. identified and cloned genes for (i) a
10 stereospecific NHase from NRRL-18668, comprising both the α -subunit of the amino acid sequence identified in the Sequence Listing by SEQ ID NO.:1 and the β -subunit of the amino acid sequence identified in the Sequence Listing by SEQ ID NO.:2; (ii) an amidase from
15 NRRL-18668 with deduced amino acid sequence identified in the Sequence Listing by SEQ ID NO.:28; (iii) a gene from NRRL-18668 designated P14K which is essential for NRRL-18668 NHase activity and with deduced amino acid sequence identified in the Sequence Listing by SEQ ID
20 NO.:22;

II. obtained DNA sequences encoding the α -subunit identified in the Sequence Listing by SEQ ID NO.:3; and the β -subunit identified in the Sequence Listing by SEQ ID NO.:4; and the amidase enzyme
25 identified in the Sequence Listing by SEQ ID NO.:27; and the P14K polypeptide identified in the Sequence Listing by SEQ ID NO.:21;

III. constructed recombinant DNA plasmids containing the genes as described in I above located
30 within an 8.0 kb DNA fragment as described in Figure 10.

IV. transformed microbial hosts with the plasmids described in III above as described in Figures 3, 15, and 16;

35 V. developed a method for the production of stereospecific NHase which comprises growing a transformed host described in IV and recovering the nitrile hydrating activity from the culture;

VI. developed a method for the production of chiral amides which comprises stereospecifically hydrating the nitrile using the nitrile hydrating activity recovered in V;

5 VII. developed a method for the production of chiral amides which comprises stereospecifically hydrating the nitrile using the nitrile hydrating activity recovered in V for the production of chiral amides using isolated microbial cells as described in
10 IV, the treated matter thereof, or a fixed form of them;

VIII. constructed recombinant DNA plasmids containing the NHase genes as described in I above, in combination with the amidase gene derived from
15 *Pseudomonas chlororaphis* B23 (FERM B-187) or the amidase gene described in I above;

IX. transformed microbial hosts with the plasmids described in VIII above as described in Figures 6 and 18;

20 X. developed a method for the production of NHase and amidase which comprises growing a transformed host described in IX and recovering the nitrile hydrating and amide hydrating activity from the culture; and

25 XI. developed a method for the production of chiral amides and chiral acids which comprises stereoselective hydration of the nitrile and its amide products using the NHase and amidase activities recovered in V for the production of the chiral
30 products using isolated microbial cells as described in IX, the treated matter thereof, or a fixed form of them to produce chiral products.

I. ISOLATION AND CLONING OF THE NITRILE HYDRATASE GENE:

35 A. Isolation and Partial Amino Acid Sequencing of the Nitrile Hydratase Enzyme:

The instant invention provides a nitrile hydratase enzyme which is defined above. The nitrile

hydratase of the present invention was isolated and purified from *Pseudomonas putida* (NRRL-18668). Bacterial nitrile hydratases are known to be generally comprised of structurally distinct α and β subunits (Hashimoto et al., *Biosci., Biotechnol., Biochem.*, 58(10), 1859-65 (1994)). The instant nitrile hydratase was separated into α and β subunits using HPLC methodology. Methods for the purification and separation of enzymes by HPLC are common and known in the art. See, for example, Rudolph et al., *Chromatogr. Sci.*, 51 (HPLC Biol. Macromol.), 333-50 (1990).

N-terminal amino acid sequences of each subunit were determined using methods well known in the art. See, for example, Matsudaira, P., *Methods Enzymol.*, 182 (Guide Protein Purif.), 602-13 (1990). Fragments of each subunit were generated and partial amino acid sequences of the fragments were determined. Partial sequences of the α and β subunits of this nitrile hydratase are shown in SEQ ID NOs.:5-9 and 10-13, respectively.

B. DNA Probe for Isolation of the Nitrile Hydratase Gene:

In order to isolate the nitrile hydratase gene, a series of degenerate 21-mer oligonucleotide primers based on the available NRRL-18668 NHase amino acid sequence were designed and synthesized for use as polymerase chain reaction (PCR) primers. Genomic DNA was isolated from *P. putida* (NRRL-18668) by standard methods (Sambrook, J., et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press (1989)) and was used as a target for PCR with numerous degenerate primer combinations. The resulting amplified products were subjected to Southern analysis (Southern, E. M., *J. Mol. Biol.*, 98, 503, (1975)) using isolated *Rhodococcus rhodochrous* J1 L-NHase gene (Kobayashi, M., *Biochem. Biophys. Acta* 1129:23-33 (1991)) as a probe. One strongly

hybridizing fragment of 0.7 kb was identified from a PCR reaction based on the degenerate primers designated D1 and D7. The sequences of D1 and D7 are identified in the Sequence Listing as SEQ ID NO.:14 and SEQ ID NO.:15, respectively. The 0.7 kb PCR fragment was subcloned into the plasmid M13 using standard methods (Sambrook, *supra*) and sequenced. Sequencing revealed that the 0.7 kb fragment demonstrated a 60% base homology to the *Rhodococcus rhodochromus* J1 L-NHase gene. Deduced amino acid sequence from this 0.7 kb fragment was compared to available NRRL-18668 amino acid sequences determined previously and to other known NHase sequences. The comparison confirmed that this fragment was part of the *P. putida* NHase gene. The 0.7 kb DNA fragment was sequenced and is identified as SEQ ID No.:16. The 0.7 kb fragment was used as a probe to isolate a genomic DNA fragment from NRRL-18668 which contains the entire NHase gene.

20 C. Isolation of a Genomic DNA Fragment
Containing NRRL-18668 NHase Gene:

Genomic DNA isolated from *P. putida* (NRRL-18668) was digested with restriction enzymes EcoRI and XhoI and size-selected by agarose gel electrophoresis based on Southern blotting using the 0.7 kb DNA fragment described above as a probe. Restricted genomic DNA was then cloned into phage lambda ZAPII [Stratagene, La Jolla, CA]. The lambda library was screened with the 0.7 kb DNA fragment probe and one positively hybridizing phage clone with a DNA insert of 6.5 kb was identified and isolated.

D. Plasmid Construction and Host Transformation
and Confirmation of NHase Sequence:

Once a positive clone containing a 6.5 kb insert was identified, the presence of the NHase gene in the clone was confirmed by a process of (i) constructing a plasmid containing the 6.5 kb insert (pSW1, Figure 1); (ii) transforming a suitable host cell with this

plasmid; (iii) growing up the transformed host and purifying the plasmid DNA; (iv) constructing a restriction map from the purified DNA (Figure 2); and (v) sequencing the NHase genes. The confirmation process is common and well known in the art and techniques used may be found in Sambrook *supra*.

Sequence analysis confirmed the nitrile hydratase coding regions, which consisted of two open reading frames corresponding to the alpha and beta subunits of the corresponding NHase protein as defined in the Sequence Listing by SEQ ID NO.:17 and Fig. 7. The α and β open reading frames were analyzed for base sequence similarly to the *Rhodococcus rhodochrous* J1 L-NHASE gene used as a probe and described above. Homology comparisons showed that the α open reading frame had 64% homology to the region encoding the α subunit on the J1 gene and the β open reading frame had 52% homology to the region encoding the β subunit on the J1 gene.

II. CONSTRUCTION OF EXPRESSION VECTOR AND EXPRESSION STRAINS:

The present invention provides a transformed host cell capable of expressing active nitrile hydratase enzyme. Generally, it is preferred if the host cell is an *E. coli*, however, it is not outside the scope of the invention to provide alternative hosts. Such alternative hosts may include, but are not limited to, members of the genera *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Saccharomyces*, *Pichia*, *Aspergillus*, *Hansenula*, and *Streptomyces*.

The present invention provides a variety of plasmids or vectors suitable for the cloning of the nitrile hydratase gene in the desired host. Suitable vectors for construction contain a selectable marker and sequences allowing autonomous replication or chromosomal integration. Additionally, suitable vectors for expression contain sequences directing transcription and translation of the heterologous DNA

fragment. These vectors comprise a region 5' of the heterologous DNA fragment which harbors transcriptional initiation controls, and optionally a region 3' of the DNA fragment which controls transcriptional termination. It is most preferred when both control regions are derived from genes homologous to the host cell, although such control regions need not be derived from the genes native to the specific species chosen as a production host. Suitable vectors can be derived, for example, from a bacteria (e.g., pET, pBR322, pUC19, pSP64, pUR278 and pORF1), a virus (such as bacteriophage T7 or a M-13 derived phage), a cosmid, a yeast or a plant. Protocols for obtaining and using such vectors are known to those in the art. (Sambrook. *supra*.)

Vectors suitable for *E. coli* will have compatible regulatory sequences and origins of replication. They will be preferably multicopy and have a selectable marker gene, for example, a gene coding for antibiotic resistance.

Promoters useful for driving the expression of heterologous DNA fragments in *E. coli* are numerous and familiar to those skilled in the art. Virtually any promoter capable of driving the gene encoding the nitrile hydratase enzyme is suitable for the present invention, although promoters native to *E. coli* are preferred and the inducible IPTG *Ptac* promoter is most preferred (deBoer, H., *Proc. Natl. Acad. Sci. USA*, 80:21-25 (1983)). Although an inducible promoter is preferred, one of skill in the art will appreciate that either inducible or constitutive promoters are suitable.

Within the context of the present invention the entire 6.5 kb DNA insert containing the NRRL-18668 NHase gene in the plasmid pSW1 was subcloned into the wide-host-range vector pMMB207 (Bagdasarian, M., *Gene*, 97:39-47 (1991)) under the control of the *Ptac* promoter to create an expression vector designated

pSW2 (Figure 3). Additionally, the 2.8 kb PstI DNA fragment derived from the 6.5 kb DNA fragment and containing the NRRL-18668 NHase gene but with substantially less upstream and downstream flanking sequence, was also subcloned into the vector pMMB207 under the control of the *Ptac* promoter to generate the plasmid pSW5 (Figure 4). Comparing these two expression constructs allowed Applicants to investigate proximal accessory sequences or proteins which might be involved in expression or activity of NHase. Applicants' studies indicated that the NHase genes may be part of an operon which generates a 10 kb mRNA transcript, of which only approximately 1.5 kb is accounted for by NHase. This suggests that additional genes are encoded by the upstream and downstream sequence flanking NHase. Others have described a requirement for downstream sequence for efficient expression of NHase in *Rhodococcus* sp. N-774 (Hashimoto, Y., *Biosci. Biotech. Biochem.*, 58:1859-1865 (1994)).

Following cloning, *E. coli* XL1-Blue host was transformed in parallel with the plasmid pSW2 or pSW5 described above. Methods of transforming host cells with foreign DNA are common and well known in the art. For example, transforming host cells with foreign DNA may be accomplished using calcium-permeabilized cells, electroporation, or by transfection using a recombinant phage virus. (Sambrook *supra*). Plasmid DNA was isolated from these transformants and enzyme restriction analysis confirmed the construction of two separate strains, one harboring the pSW2 plasmid and the other harboring the pSW5 plasmid.

The gene encoding the α subunit, and the gene encoding the β subunit of NRRL-18668 NHase were also expressed in an alternative host, the methylotrophic yeast *Pichia pastoris*. Methods for producing heterologous proteins in *P. pastoris* are well known in the art. For each subunit, the coding sequence was

placed under control of the methanol inducible promoter, alcohol oxidase I (AOX1), in a vector which was subsequently integrated into the host chromosome. Each subunit was produced in the respective host after
5 induction by methanol. NHase activity was not reproducibly obtained upon mixing extract prepared from the α producing strain with extract prepared from the β producing strain. In addition, a single strain producing both α and β subunits under control of the
10 AOX1 promoter was constructed. Both subunits were produced in this recombinant *P. pastoris* strain, but NHase activity was not obtained.

Applicants sequenced DNA both upstream and downstream of the NHase genes, and identified at least
15 two open reading frames, one upstream and one downstream. The upstream open reading frame was determined to encode an amidase enzyme, based on comparison of the deduced amino acid sequence to other amidase amino acid sequences. Plasmids were
20 constructed for the expression of NRRL-18668 amidase in *E. coli*. A search of the protein database with the deduced amino acid sequence encoded by the downstream open reading frame (designated P14K) indicated no significant matches. Plasmids were constructed for
25 expression of NHase genes only or NHase and P14K genes in both *E. coli* and *P. pastoris*. In both *E. coli* and *P. pastoris*, NHase activity was obtained only when P14K was co-expressed with the NHase genes. The preference for hydrolysis of S-nitriles (stereo-
30 specificity) observed in the native organism was also demonstrated in the recombinant organisms producing active NHase.

III. EXPRESSION OF THE NITRILE HYDRATASE ENZYME AND CONVERSION OF SUBSTRATES:

35 Transformed *E. coli* cells harboring plasmid pSW2 under the control of the IPTG inducible *Ptac* promoter, were grown under standard conditions and induced to express the nitrile hydratase enzyme. Cells were

harvested and lysed and the protein was detected in crude lysates by SDS-polyacrylamide gel electrophoresis followed by western blot analysis (Egger et al., *Mol. Biotechnol.*, 1(3), 289-305 (1994)) using antisera raised against NRRL-18668 NHase protein (Figure 5). Under these conditions induced cells produced approximately 10-fold as much nitrile hydratase protein as uninduced cells. Nitrile hydratase was not detected from a control strain harboring the vector pMMB207 without the 6.5 kb insert.

Nitrile hydratase is typically confirmed by incubating a suitable substrate nitrile in the presence of the crude or purified enzyme. Suitable substrates for the instant hydratase include a variety of racemic alkyl nitriles such as methacrylonitrile, methylbutyronitrile and propionitrile. In the instant case, nitrile hydratase activity was confirmed by monitoring the conversion of methacrylonitrile to the corresponding amide. Induced cells harboring the plasmid pSW2 showed rapid conversion of methacrylonitrile, while induced cells without the pSW2 plasmid showed no conversion of methacrylonitrile. Additionally, induced cells harboring the plasmid pSW5 show no conversion of methacrylonitrile.

Stereospecific activity of the enzyme produced in induced cells harboring plasmid pSW2 was confirmed by monitoring the conversion of R,S-CFIN to amide products using reverse-phase or chiral high pressure liquid chromatography (HPLC). Methods of enantiomer separation on HPLC are well known in the art. See, for example, Mutton, I., *Pract. Approach Chiral Sep., Liq. Chromatogr.*, 329-55 (1994), Editor(s): Subramanian, Ganapathy, Publisher: VCH, Weinheim, Germany.

IV. CO-EXPRESSION OF NITRILE HYDRATASE AND AMIDASE:

The present invention further provides a transformed microorganism capable of co-expressing

both a heterologous nitrile hydratase gene and a heterologous amidase gene. This transformant is capable of effecting the conversion of racemic mixtures of aryl-2-alkane nitriles to the
5 corresponding carboxylic acids via the amide intermediate.

A number of amidase encoding genes may be suitable for co-expression with the instant nitrile hydratase. However, the amidase gene isolated from
10 *Pseudomonas chlororaphis* B23 and defined above is preferred.

The gene encoding the *Pseudomonas chlororaphis* B23 amidase is known (Nishiyama, M. J., *Bacteriol.*, 173:2465-2472 (1991)) and was obtained through PCR
15 amplification using appropriate primers. The amplified gene comprising 1.5 kb was subcloned into a pMMB207 plasmid (already containing the nitrile hydratase gene) using standard restriction enzyme digestion and ligation techniques (Sambrook *supra*) to
20 generate the plasmid pSW17 (Figure 6). The plasmid pSW17 was constructed so as to place the amidase gene and the nitrile hydratase gene both under the control of the same IPTG inducible *Ptac* promoter. The plasmid pSW17 was then used to transform a suitable host cell
25 (e.g., *E. coli* XL1-Blue) according to standard methods.

In order to confirm the activity of the amidase produced in cells transformed with plasmid pSW17, cells transformed by plasmid pSW17 were grown up and
30 induced with IPTG in the presence of a suitable nitrile and the chiral amide and free acid products were identified by chiral HPLC analysis.

The following Examples are meant to illustrate the invention but should not be construed as limiting
35 it in any way.

EXAMPLE 1ISOLATION, PURIFICATION, AND AMINO ACID SEQUENCING
OF PORTIONS OF THE NITRILE HYDRATASE α AND β SUBUNITS

Pseudomonas putida (NRRL-18668) was cultured in a
5 medium (10 g/L glucose, 8.7 g/L K_2HPO_4 , 6.8 g/L KH_2PO_4 ,
2.0 g/L acetonitrile, 1.85 g/L $NaNO_3$, 0.50 g/L
 $MgSO_4 \cdot 7H_2O$, 0.050 g/L $FeSO_4 \cdot 7H_2O$, 0.30 mg/L $MnCl_2 \cdot 4H_2O$,
0.10 mg/L H_3BO_3 , 0.050 mg/L $NiSO_4 \cdot 6H_2O$, 0.050 mg/L
 $CuSO_4 \cdot 5H_2O$, 0.050 mg/L $Co(NO_3)_2 \cdot 6H_2O$, 0.030 mg/L
10 $Na_2MoO_4 \cdot 2H_2O$, 0.030 mg/L $ZnSO_4 \cdot 4H_2O$, 0.020 mg/L KI,
0.020 mg/L KBr, 0.010 mg/L pyridoxine-HCl, 0.0050 mg/L
thiamine-HCl, 0.0050 mg/L D-pantothenate, Ca^{2+} salt,
0.0050 mg/L riboflavin, 0.0050 mg/L nicotinic acid,
0.0050 mg/L p-aminobenzoic acid, 0.0020 mg/L biotin,
15 0.0020 mg/L vitamin B_{12} , 0.0020 mg/L folic acid,
pH 7.0) at 30°C for 48 h. The bacterial cells were
harvested. 100 g of the bacterial cells were
disrupted and the cell free extract fractionated with
ammonium sulfate. The ammonium sulfate fractionation
20 precipitate was dissolved in buffer and loaded on a
Phenyl Sepharose CL-4B chromatography column
(Pharmacia Biotech, Uppsala, Sweden), followed by a
DEAE-cellulose chromatography column, and a second
DEAE-cellulose chromatography column (Whatman,
25 Maidstone, England). Active fractions were pooled and
concentrated. The concentrate containing the enzyme
was loaded on a reverse phase high performance
chromatography column (Vydac 208TP104) and two
subunits (α and β) were obtained. The N-terminal
30 amino acid sequence of the α - and β -subunits was
determined using an amino acid sequencer (Beckman
model LF3000G gas phase protein sequencer, Fullerton,
CA. The α - and β -subunits were cleaved separately
using cyanogen bromide, TPCK-treated trypsin, and AspN
35 protease, and the peptides generated were separated on
a reverse phase high performance chromatography column
(Vydac 208TP104, The Separations Group, Hesperia, CA).
Fractions containing well-resolved peptides were

sequenced using the same technique. The sequences of the individual peptides were combined into partial sequences of the subunits by alignment with the published sequences of the α - and β -subunits of nitrile hydratases from *P. chlororaphis* B23 [Nishiyama et al., *J. Bacteriol.*, 173:2465-2472 (1991)], *Rhodococcus* N-774 [Ikehata et al., *Eur. J. Biochem.*, 181:563-570 (1989)], and *Rhodococcus rhodochrous* J1 [Kobayashi et al., *Biochim. Biophys. Acta*, 1129:23-33 (1991)]. The partial sequences of the α - and β -subunits of nitrile hydratase from *Pseudomonas putida* (NRRL-18668) were identified as defined in the Sequences Listing as SEQ ID NOS.:5-9 and SEQ ID NOS.:10-13, respectively.

EXAMPLE 2

PREPARATION OF DNA PROBE FOR NRRL-18668 NHASE GENE

The degenerate oligonucleotide designated D1 as defined in the Sequence Listing as SEQ ID NO.:14, and the degenerate oligonucleotide designated D7 as defined in the Sequence Listing as SEQ ID NO.:15 were used as primers in a polymerase chain reaction (PCR) [Mullis, K. B., *Meth. Enzymol.*, 155:335-350 (1987)] with NRRL-18668 genomic DNA as target. PCR conditions were as follows: 100 ng target, 1 μ M each primer, 200 μ M each of dATP, dCTP, dGTP, dTTP, 10 mM Tris-HCl pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.001% gelatin, 25 U/mL *Amplitaq*[™] DNA polymerase (Perkin Elmer Cetus, Norwalk, CT). PCR parameters were as follows: 94°C 1 min, 55°C 1 min, 72°C 1 min, 40 cycles. One half of the PCR product was subjected to ethidium bromide agarose gel electrophoresis followed by transfer to nitrocellulose and Southern analysis with ³²P labeled *Rhodococcus rhodochrous* J1 L-NHase gene as probe [Southern, E.M., *J. Mol. Biol.*, 98:503 (1975)]. Strong hybridization of a DNA fragment of approximately 0.7 kb suggested the presence of at least a portion of a NHase gene in this PCR product. The remaining half of the PCR product was restricted

with EcoR1 (the primers were designed with EcoR1 sites at the 5' ends) and ligated to EcoR1 restricted M13 mp19 vector DNA. Ligation mix was used to transfect competent *E. coli* XL1-Blue which was plated onto LB plates supplemented with IPTG and X-gal (5-bromo-4chloro-3indolyl- β -D-galactopyranoside) [Maniatis, T., Molecular Cloning: A Laboratory Manual (1989)]. Phage DNA was prepared from several "white" plaques [Maniatis, T., Molecular Cloning: A Laboratory Manual (1989)] and sequenced by dideoxy termination protocol using universal primer [Sanger, F., Science, 214:1205-1210 (1981)]. Analysis of the nucleotide sequence obtained as defined in the Sequence Listing as SEQ ID NO.:16 confirmed that the PCR product corresponds to part of the NHase gene.

EXAMPLE 3

ISOLATION OF GENOMIC DNA FRAGMENT CONTAINING NRRL-18668 NEASE GENE

Total genomic DNA (10 μ g) from NRRL-18668 was isolated [Maniatis, T., Molecular Cloning: A Laboratory Manual (1989)], restricted with EcoR1 and Xho1, and one half subjected to agarose gel electrophoresis followed by Southern blot using the 32 P labeled 0.7 kb fragment described in Example 2 as a probe [Southern, E. M., *J. Mol. Biol.*, 98:503 (1975)]. A strongly hybridizing band of approximately 6.5 kb was identified, suggesting that the NHase gene (or part of it) resides on this 6.5 kb genomic DNA fragment. A duplicate agarose gel was run and a gel slice from the 6.5 kb region was excised. DNA extracted from the gel slice isolated [Maniatis, T., Molecular Cloning: A Laboratory Manual (1989)] was ligated to lambda DNA restricted with EcoR1 and Xho1. The ligation mix was packaged into phage particles and used to transfect *E. coli* XL1-Blue according to the manufacturer's instructions [Stratagene, La Jolla, CA]. Several thousand plaques were screened using the 32 P-labeled 0.7 kb fragment as probe [Maniatis, T.,

Molecular Cloning: A Laboratory Manual (1989)]. One positively hybridizing plaque was subsequently purified.

EXAMPLE 4

5 CONSTRUCTION OF PLASMID CONTAINING NRRL-18668 NHASE GENE

DNA from the purified phage plaque described in Example 3 was excised and converted to a pBluescript-based plasmid according to the manufacturer's
10 instructions [Stratagene, La Jolla, CA], and designated pSW1. The plasmid pSW1 has a 6.5 kb insert containing the NRRL-18668 NHase gene as described in Fig. 1.

EXAMPLE 5

15 TRANSFORMATION OF HOST BY PLASMID CONTAINING NRRL-18668 NHASE GENE

The plasmid pSW1 described in Example 4 was used to transform competent *E. coli* XL1-Blue cells by the CaCl_2 method [Maniatis, T., Molecular Cloning: A
20 Laboratory Manual (1989)].

EXAMPLE 6

RECOMBINANT PLASMID PURIFICATION AND CONSTRUCTION OF RESTRICTION MAP FOR 25 GENOMIC DNA FRAGMENT CONTAINING NRRL-18668 NHASE GENE

Plasmid DNA purified by the alkaline lysis method [Maniatis, T., Molecular Cloning: A Laboratory Manual (1989)] from *E. coli* cells harboring plasmid pSW1, described in Example 5, was restricted with EcoRI, PstI, KpnI, Hind3, and XhoI singly or in various
30 combinations, followed by agarose gel analysis, and Southern analysis using the 0.7 kb PCR product described in Example 2 as a probe [Southern, E.M., J. Mol. Biol., 98:503 (1975)]. A restriction map
35 constructed for the 6.5 kb insert fragment of the plasmid pSW1, including the location of the NHase gene is shown in Fig. 2.

EXAMPLE 7DNA SEQUENCING OF NRRL-18668 NHASE GENE

Based on the restriction map described in Example 6, the nucleotide sequence of a fragment of DNA encompassing the NHase gene was determined by the Sanger dideoxy method [Sanger, F., Science, 214:1205-1210 (1981)] using double-stranded plasmid DNA as template. The nucleotide sequence and the corresponding predicted amino acid sequences for the α and β peptides are defined in the Sequence Listing as SEQ ID NO.:17 and Fig. 7.

EXAMPLE 8CONSTRUCTION OF NRRL-18668 NHASE EXPRESSION VECTOR

Plasmid pSW1 was restricted with EcoR1 and Xho1 and the 6.5 kb fragment was ligated to the wide host range plasmid pMMB207 [Bagdasarian, M., Gene, 97:39-47 (1991)] restricted with EcoR1 and Sal 1 to generate the plasmid designated pSW2 and shown in Fig. 3. The 2.8 kb Pst1 DNA fragment containing the NRRL-18668 NHase gene was excised from plasmid pSW2 by digestion with Pst1 restriction enzyme and ligated into the Pst1 site of vector pMMB207 to generate the plasmid designated pSW5 and shown in Fig. 4.

EXAMPLE 9CONSTRUCTION OF NRRL-18668 NHASE EXPRESSION STRAIN

Plasmids pSW2 and pSW5 described in Example 8 were used to transform competent *E. coli* XL1-Blue cells which were plated onto LB plates supplemented with 12.5 μ g/mL chloramphenicol [Maniatis, T., Molecular Cloning: A Laboratory Manual (1989)].

EXAMPLE 10EXPRESSION OF NRRL-18668 NHASE PROTEIN

E. coli cells harboring plasmid pSW2, described in Example 8A, were grown in SOC media (0.5 g/L NaCl, 20 g/L bacto-tryptone, 5 g/L bacto-yeast extract, 20 mM glucose, 2.5 mM KCl, 10 mM MgCl₂) at 37°C to OD₆₀₀=0.5, followed by induction at 30°C by the addition of IPTG to 1 mM. After induction times

ranging from 0.5 h to 3 h, cells were harvested by centrifugation, and suspended in 1/10 volume PBS (8.0 g/L NaCl, 0.2 g/L KCl, 1.44 g/L Na₂HPO₄, 0.24 g/L KH₂PO₄ pH 7.4). A cell suspension equivalent to 0.05 OD600 units is added to an equal volume of 2X SDS gel-loading buffer (100 mM Tris pH 6.8, 200 mM DTT, 4% SDS, 0.2% bromophenol blue, 20% glycerol), boiled for 5 min, and analyzed by SDS PAGE [Laemmli, U.K., *Nature*, 227:680-685 (1970)] followed by western blot [Towbin, H., *Proc. Natl. Acad. Sci.* 76:4350-4354 (1979)] using antisera raised against NRRL-18668 NHase protein. A positive signal was obtained at approximately 28 kd and corresponded to purified NHase protein as shown in Fig. 5.

15

EXAMPLE 11

EXPRESSION OF ACTIVE NRRL-18668 NHASE

E. coli cells harboring plasmid pSW2, described in Example 9, were grown and induced as described in Example 9 in a 500 mL batch. Cells were harvested by centrifugation and washed with pH 7.2, 0.1M phosphate buffer (KH₂PO₄ adjusted with 50% NaOH) containing 15% glycerol. Washed cells were stored frozen at -70°C. Washed and frozen *E. coli* cells harboring the pSW2 plasmid and were suspended in 100 mM phosphate buffer, pH 7, at a cell density of O.D.₄₉₀ = 0.62. Methacrylonitrile was added to a final concentration of 10 mM and the mixture was shaken at 250 rpm at room temperature. Analysis of supernatant showed that methacrylonitrile was rapidly converted to hydrolysis products after 30 min. Cells without the pSW2 plasmid showed no activity.

25

30

EXAMPLE 12

PRODUCTION OF CHIRAL AMIDES

Induced *E. coli* cells harboring the pSW2 plasmid and producing stereospecific nitrile hydratase activity as described in Example 11 were suspended in 100 mM phosphate buffer, pH 7, and a concentration of 50 mg/mL. One milliliter of this suspension was

35

placed in a glass vial containing 19.3 mg of R,S-CPIN. The suspension was shaken at 250 rpm on a rotary shaker at room temperature for 68 h. Analysis by chiral HPLC reveals only the S-CPIAm was produced from the R,S-CPIN.

| Time, h | mg nitrile | | mg amide | |
|---------|------------|--------|----------|---------|
| | R-CPIN | S-CPIN | R-CPIAm | S-CPIAm |
| 0 | 9.6 | 9.6 | 0 | 0 |
| 68 | 9.6 | 5.5 | 0 | 4.5 |

EXAMPLE 13

CONSTRUCTION OF A VECTOR FOR CO-EXPRESSION OF NRRL-18668 NHASE AND PSEUDOMONAS CHLORORAPHIS B23 AMIDASE

The amidase gene from *Pseudomonas chlororaphis* B23 (defined as SEQ ID NO.:20) was obtained through PCR amplification using primers with overhanging 5' EcoR1 sites as defined in the Sequence Listing as SEQ ID NO.:18 and SEQ ID NO.:19. This 1.4 kb DNA fragment containing the B23 amidase gene was digested with EcoR1 restriction enzyme and ligated into the EcoR1 site of pMMB207, and the 5.0 kb EcoR1/Hind111 DNA fragment from pSW1, described in Example 4, was subcloned between the Xba1 and Hind111 to generate the plasmid pSW17 as shown in Fig. 6.

EXAMPLE 14

CONSTRUCTION OF STRAIN FOR CO-EXPRESSION OF NRRL-18668 NHASE AND PSEUDOMONAS CHLORORAPHIS B23 AMIDASE

Plasmid pSW17 described in Example 13 was used to transform competent *E. coli* XL1-Blue cells which were selectively grown on LB plates supplemented with 12.5 µg/mL chloramphenicol [Maniatis, T., Molecular Cloning: A Laboratory Manual (1989)].

EXAMPLE 15

COMPARISON OF NHase ACTIVITY FROM pSW2 AND pSW5

E. coli cells harboring the pSW2 or pSW5 plasmid and induced according to the protocol in Example 11 were each suspended separately in 100 mM phosphate buffer, pH 7, at a concentration of 20 mg/mL.

Butyronitrile was added to each suspension to a final concentration of 10 mM. The suspensions were shaken at 250 rpm on a rotary shaker at room temperature for 24 h. At the end of the incubation period, 0.1% phosphoric acid was added to the suspensions, bringing them to a pH of 2-3 and stopping nitrile hydratase activity. Cells were removed from the suspension by centrifugation. Analysis of the reactions showed the following products:

- 10 pSW2 - 94% butyramide, 6% butyronitrile;
 pSW5 - <1% butyramide, 100% butyronitrile.

EXAMPLE 16

PRODUCTION OF S-CPIAM AND S-CPIA FROM R,S-CPIN

E. coli cells harboring the pSW17 and induced according to the protocol in Example 11 were suspended in 100 mM phosphate buffer, pH 7, at a concentration of 100 mg/mL. One milliliter of this suspension was placed in a glass vial containing 19.3 mg of R,S-CPIN dispersed in a dry form on 0.5 g of 0.5 mm glass beads. The suspension was shaken in a 20 mL scintillation vial at 250 rpm on a rotary shaker at room temperature for 68 h. Analysis by chiral HPLC reveals both S-CPIAm and the S-CPIA were produced from the R,S-CPIN.

| Time, h | mg nitrile | | mg amide | | mg acid | |
|---------|------------|--------|----------|---------|---------|--------|
| | R-CPIN | S-CPIN | R-CPIAm | S-CPIAm | R-CPIA | S-CPIA |
| 0 | 9.6 | 9.6 | 0 | 0 | 0 | 0 |
| 68 | 9.6 | 8.4 | 0 | 0.84 | 0 | 0.42 |

25

EXAMPLE 17

Nucleotide sequencing of DNA regions flanking NRRL-18668 NHase gene

The nucleotide sequences of DNA regions flanking the NRRL-18668 NHase were determined by the Sanger dideoxy method (Sanger, F. (1981) Science 214:1205-1210) using double-stranded plasmid DNA as template. Using pSW1 (Fig. 1) as template, the nucleotide sequence downstream of NHase, down to the

XhoI site (Fig. 2), was determined. This sequence contains at least one gene, and potentially several more, which is defined as P14K, the nucleotide sequence of which is defined in Sequence Listing SEQ ID NO.:21, and the deduced amino acid sequence is defined in Sequence Listing SEQ ID NO.:22. P14K is required for NHase activity as described below.

The nucleotide sequence upstream of NHase, up to the EcoRI (Fig. 2), was determined using pSW1 (Fig. 1) as template. The nucleotide sequence further upstream of the EcoRI site was determined after subcloning DNA fragments corresponding to this region as follows. NRRL-18668 genomic DNA was digested with PstI and then self-ligated. Oligo-nucleotide primers designed to bind 3' to EcoRI heading upstream (Fig. 2) and 5' to PstI heading downstream (Fig. 2), and defined as Sequence Listing SEQ ID NO.:23 and Sequence Listing SEQ ID NO.:24, respectively, were used in a PCR reaction to amplify a 0.8 kb fragment corresponding to DNA upstream of the EcoRI site (Fig. 8). NRRL-18668 genomic DNA was digested with EcoRI and then self-ligated. Oligo-nucleotide primers designed to bind 3' to PstI heading upstream (Fig. 8) and 5' to EcoRI heading downstream (Fig. 8), and defined as Sequence Listing SEQ ID NO.:25 and SEQ ID NO.:26, respectively, were used in a PCR reaction to amplify a 0.7 kb fragment corresponding to DNA upstream of the PstI site (Fig. 9). By subcloning and sequencing the PCR fragments, the nucleotide sequence upstream of NHase, up to the EcoRI site (Fig. 9) was determined. This sequence contains at least one gene, and potentially more, which has been identified as encoding an amidase (based on homology to other amidase sequences), the nucleotide sequence of which is defined as Sequence Listing SEQ ID NO.:27, and the deduced amino acid sequence defined as Sequence Listing SEQ ID NO.:28.

A compiled map of the entire 8.0 kb DNA fragment, indicating genes identified, is shown in Fig. 10.

EXAMPLE 18Construction of plasmids for expression
of NRRL-18668 NHase in *Pichia pastoris*

The 0.9 kb EcoR1/Xba1 fragment in pHIL-D4
5 (Phillips Petroleum, Bartlesville, OK) was replaced by
the 0.9 kb EcoR1/Xba1 fragment from pAO815
(Invitrogen, San Diego, CA) to generate the plasmid
pHIL-D4B2 (Fig. 11) which contains the following
elements: 5'AOX1, *P. pastoris* methanol inducible
10 alcohol oxidase I (AOX1) promoter; AOX1 term,
P. pastoris AOX I transcriptional termination region;
HIS4, *P. pastoris* histidinol dehydrogenase-encoding
gene for selection in *his4* hosts; kan, sequence
derived from transposon Tn903 encoding aminoglycoside
15 3'-phosphotransferase, conferring kanamycin, neomycin
and G418 resistance in a wide variety of hosts, and
useful as an indicator of cassette copy number;
3'AOX1, *P. pastoris* sequence downstream from AOX1,
used in conjunction with 5'AOX1 for site-directed
20 vector integration; ori, pBR322 origin of DNA
replication allowing plasmid manipulations in *E. coli*;
and amp, β -lactamase gene from pBR322 conferring
resistance to ampicillin. An additional feature of
pHIL-D4B2 is that multiple expression cassettes
25 (5'AOX1 - gene - AOX1term) can easily be placed into
one plasmid by subcloning cassettes on Bgl2/Xba1
fragments into BamH1/Xba1 sites.

The genes encoding α , β , and P14K (Fig. 10) were
PCR amplified using primers with EcoR1 sites at the 5'
30 ends. The PCR products were digested with EcoR1, and
subcloned into the EcoR1 site of pHIL-D4B2 to generate
pSW46 (Fig. 12), pSW47 (Fig. 13) and pSW48 (Fig. 14),
respectively. The Bgl2/Xba1 fragment from pSW47
containing the β expression cassette was subcloned into
35 the BamH1/Xba1 sites of pSW46 to generate pSW49
(Fig. 15), which contains expression cassettes for α
and β . The Bgl2/Xba1 fragment from pSW48 containing
the P14K expression cassette was subcloned into the

BamHI/XbaI sites of pSW49 to generate pSW50 (Fig. 16), which contains expression cassettes for α , β and P14K.

EXAMPLE 19

Construction of *Pichia pastoris* strain for expression of NRRL-18668 NHase

5 *P. pastoris* strain GTS115(*his4*) (Phillips Petroleum, Bartlesville, OK) was transformed with 1-2 μ g of BglII-linearized plasmid pSW49 or 1-2 μ g of BglII-linearized plasmid pSW50 using the spheroplast transformation method as described (Cregg et al. 10 (1985) *Mol. Cell. Biol.* 5: 3376-3385). Cells were regenerated on plates without histidine for 3-4 d at 30°C. All transformants arise after integration of plasmid DNA into the chromosome. Chromosomal DNA was 15 prepared from *his*⁺ transformants and subjected to PCR analysis with primers specific for α , β and P14K genes. An isolated pSW49 transformant positive for α and β genes, and an isolated pSW50 transformant positive for α , β and P14K genes, designated SW49 and 20 SW50.2, respectively, were selected for further study. *P. pastoris* strain SW50.2 was deposited with ATCC and assigned accession number ATCC 74391.

EXAMPLE 20

NRRL-18668 NHase activity in engineered *P. pastoris*

25 *P. pastoris* strains SW49 and SW50.2 were grown to A_{600} of 2-10 in MGY (1.34% yeast nitrogen base without amino acids, 0.00004% biotin, 1% glycerol) with shaking at 30 °C. Cells are then pelleted and induced by resuspending in MM (1.34% yeast nitrogen base 30 without amino acids, 0.00004% biotin, 0.5% methanol) and incubated with shaking at 30 °C for 1-4 d. Cells were harvested by centrifugation and washed in PBS (0.1 M KH_2PO_4 , pH 7.2). NHase activity was demonstrated by methacrylonitrile assay, in which 35 cells were resuspended in PBS at A_{600} of 0.6, and methacrylonitrile was added to a final concentration of 10 mM. After incubation with shaking at room temperature, conversion of methacrylonitrile to

methacrylamide by NHase was demonstrated by monitoring the increase in A_{224} of the supernatant. Cells boiled before assay serve as a negative control. NHase activity was observed in SW50.2 which harbors expression cassettes for α , β and F14K, while SW49, which only harbors expression cassettes for α and β showed negligible NHase activity.

| <u>rxn time, min</u> | A_{224} | | |
|----------------------|-------------|---------------|--------------------|
| | <u>SW49</u> | <u>SW50.2</u> | <u>SW50.2 boil</u> |
| 0 | 0.260 | 0.360 | 0.110 |
| 15 | 0.360 | 1.390 | 0.125 |

Stereospecific NHase activity was also demonstrated in induced SW50.2 cells by using R-2-(4-chlorophenyl)-3-methylbutyronitrile (R-CPIN) or S-2-(4-chlorophenyl)-3-methylbutyronitrile (S-CPIN) as substrate and then analyzing for conversion to the corresponding amides (R-CPIAm and S-CPIAm, respectively) by HPLC.

| <u>rxn time, h</u> | mM | | | |
|--------------------|---------------|----------------|---------------|----------------|
| | <u>R-CPIN</u> | <u>R-CPIAm</u> | <u>S-CPIN</u> | <u>S-CPIAm</u> |
| 0 | 10 | 0 | 10 | 0 |
| 48 | 10 | 0 | 5.5 | 4.5 |

Bioconversion of adiponitrile (ADN) to 5-cyanovaleramide (5-CVAm) was also demonstrated in permeabilized SW50.2 cells, and in SW50.2 cell extracts. Permeabilized cells were prepared by the addition of benzalkonium chloride (Lonza Baequat MB-50) to a 10% (wt) suspension of induced cells to yield 1% (wt MB-50:wt cells). The suspension was then mixed on a nutator mixer for 60 min at room temperature, after which cells were washed by centrifugation 3 times with 50 mM phosphate buffer, pH 7.0. Extracts were prepared by rapidly vortexing induced cells with 0.5 mm glass beads (BioSpec Products) in 50 mM KH_2PO_4 , pH 7.0/1 mM EDTA/0.1 mM PMSF for 2 min. NHase activity was determined to be

34-38 U/g wet wt (permeabilized cells), and 35-56 U/g wet wt (cell extracts).

EXAMPLE 21

Construction of plasmid for expression of NRRL-18668 amidase in *E. coli*

5 The gene encoding NRRL-18688 amidase was PCR amplified using an upstream primer with a Hind3 site at the 5' end and a downstream primer with an Xho1 site at the 5' end. The PCR product was subcloned
10 into the vector pET-21a(+) (Novagen, Madison, WI) between the Hind3 and Xho1 sites to generate the expression plasmid pSW37 (Fig. 17).

EXAMPLE 22

Construction of *E. coli* strain for expression of NRRL-18668 amidase

15 *E. coli* strain BL21(DE3) (Novagen, Madison, WI) was transformed with pSW37 using the calcium chloride procedure (Maniatis et al. (1989) Molecular Cloning: A Laboratory Manual), and an isolated transformant was
20 designated SW37, and deposited with ATCC and assigned accession number ATCC 98174. Induced SW37 shows production of amidase enzyme based on Coomassie Blue stained denaturing polyacrylamide gel electrophoresis of soluble cell extract.

EXAMPLE 23

NRRL-18668 amidase activity in engineered *E. coli*

25 *E. coli* strain SW37 is grown in LB media at 30 °C to A₆₀₀=0.5, at which time IPTG is added to 1 mM and incubation continued for 2 h. Cells are then pelleted
30 and washed in PBS. Cells are incubated with 10 mM butyramide and conversion to butyric acid is monitored by HPLC.

EXAMPLE 24

Construction of plasmid for expression of NRRL-18668 amidase and NHase in *E. coli*

35 The entire 8.0 kb DNA fragment (shown in Fig. 10) was subcloned between the EcoR1 and Xho1 sites of the

vector pET-21(+) (Novagen, Madison, WI) to generate the plasmid pSW23 (Fig. 18).

EXAMPLE 25

Construction of *E. coli* strain for co-expression of NRRL-18668 amidase and NHase

E. coli strain BL21(DE3) (Novagen, Madison, WI) was transformed with pSW23 using the calcium chloride procedure (Maniatis et al. (1989) Molecular Cloning: A Laboratory Manual), and an isolated transformant was designated SW23, and deposited with ATCC and assigned accession number ATCC 98175. Induced SW23 shows production of NHase enzyme and amidase enzyme based on Coomassie Blue stained denaturing polyacrylamide gel electrophoresis of soluble cell extract.

EXAMPLE 26

NRRL-18668 amidase and NHase activity in engineered *E. coli*

E. coli strain SW23 is grown in LB media at 30 °C to $A_{600}=0.5$, at which time IPTG is added to 1 mM and incubation continued for 2 h. Cells are then pelleted and washed in PBS. Cells are incubated with 10 mM butyronitrile and conversion to butyric acid is monitored by HPLC. Stereospecific conversion of S-CPIN, relative to R-CPIN, to the corresponding acid (S-CPIAc) can also be monitored by HPLC.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT:
 - (A) NAME: E. I. DU PONT DE NEMOURS AND COMPANY
 - (B) STREET: 1007 MARKET STREET
 - (C) CITY: WILMINGTON
 - (D) STATE: DELAWARE
 - (E) COUNTRY: UNITED STATES OF AMERICA
 - (F) POSTAL CODE (ZIP): 19898
 - (G) TELEPHONE: 302-892-8112
 - (H) TELEFAX: 302-773-0164
 - (I) TELEX: 6717325
- (ii) TITLE OF INVENTION: NUCLEIC ACID FRAGMENTS
ENCODING STEREOSPECIFIC
NITRILE HYDRATASE AND
AMIDASE ENZYMES AND
RECOMBINANT ORGANISMS
EXPRESSING THOSE ENZYMES
USEFUL FOR THE
PRODUCTION OF CHIRAL
AMIDES AND ACIDS
- (iii) NUMBER OF SEQUENCES: 28
- (iv) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: FLOPPY DISK
 - (B) COMPUTER: IBM PC COMPATIBLE
 - (C) OPERATING SYSTEM: MICROSOFT WINDOWS 3.1
 - (D) SOFTWARE: MICROSOFT WORD 2.0C
- (v) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vi) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 60/004914
 - (B) FILING DATE: OCTOBER 5, 1995
- (vii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: FLOYD, LINDA A.
 - (B) REGISTRATION NO.: 33,692
 - (C) REFERENCE/DOCKET NUMBER: CR-9677-A

(2) INFORMATION FOR SEQ ID NO.:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 210 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: unknown
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:1:

```

Met Gly Gln Ser His Thr His Asp His His His Asp Gly Tyr Gln Ala
1      5      10      15
Pro Pro Glu Asp Ile Ala Leu Arg Val Lys Ala Leu Glu Ser Leu Leu
20     25     30
Ile Glu Lys Gly Leu Val Asp Pro Ala Ala Met Asp Leu Val Val Gln
35     40     45
Thr Tyr Glu His Lys Val Gly Pro Arg Asn Gly Ala Lys Val Val Ala
50     55     60
Lys Ala Trp Val Asp Pro Ala Tyr Lys Ala Arg Leu Leu Ala Asp Ala
65     70     75     80
Thr Ala Ala Ile Ala Glu Leu Gly Phe Ser Gly Val Gln Gly Glu Asp
85     90     95
Met Val Ile Leu Glu Asn Thr Pro Ala Val His Asn Val Phe Val Cys
100    105    110
Thr Leu Cys Ser Cys Tyr Pro Trp Pro Thr Leu Gly Leu Pro Pro Ala
115    120    125
Trp Tyr Lys Ala Ala Ala Tyr Arg Ser Arg Met Val Ser Asp Pro Arg
130    135    140
Gly Val Leu Ala Glu Phe Gly Leu Val Ile Pro Ala Asn Lys Glu Ile
145    150    155    160
Arg Val Trp Asp Thr Thr Ala Glu Leu Arg Tyr Met Val Leu Pro Glu
165    170    175
Arg Pro Gly Thr Glu Ala Tyr Ser Glu Glu Gln Leu Ala Glu Leu Val
180    185    190
Thr Arg Asp Ser Met Ile Gly Thr Gly Leu Pro Thr Gln Pro Thr Pro
195    200    205
Ser His
210

```

(2) INFORMATION FOR SEQ ID NO.:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 217 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: unknown
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:2:

```

Met Asn Gly Ile His Asp Thr Gly Gly Ala His Gly Tyr Gly Pro Val
1      5      10      15
Tyr Arg Glu Pro Asn Glu Pro Val Phe Arg Tyr Asp Trp Glu Lys Thr
20      25      30
Val Met Ser Leu Leu Pro Ala Leu Leu Ala Asn Ala Asn Phe Asn Leu
35      40      45
Asp Glu Phe Arg His Ser Ile Glu Arg Met Gly Pro Ala His Tyr Leu
50      55      60
Glu Gly Thr Tyr Tyr Glu His Trp Leu His Val Phe Glu Asn Leu Leu
65      70      75      80
Val Glu Lys Gly Val Leu Thr Ala Thr Glu Val Ala Thr Gly Lys Ala
85      90      95
Ala Ser Gly Lys Thr Ala Thr Arg Val Leu Thr Pro Ala Ile Val Asp
100     105     110
Asp Ser Ser Ala Pro Gly Leu Leu Arg Pro Gly Gly Gly Phe Ser Phe
115     120     125
Phe Pro Val Gly Asp Lys Val Arg Val Leu Asn Lys Asn Pro Val Gly
130     135     140
His Thr Arg Met Pro Arg Tyr Thr Arg Ala Lys Trp Gly Gln Trp Ser
145     150     155     160
Ser Thr Met Val Cys Phe Val Thr Pro Asp Thr Ala Ala His Gly Lys
165     170     175
Gly Glu Gln Pro Gln His Val Tyr Thr Val Ser Phe Thr Ser Val Glu
180     185     190
Leu Trp Gly Gln Asp Ala Ser Ser Pro Lys Asp Thr Ile Arg Val Asp
195     200     205
Leu Trp Asp Asp Tyr Leu Glu Pro Ala
210     215

```

(2) INFORMATION FOR SEQ ID NO.:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 633 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:3:

```

ATGGGGCAAT CACACACGCA TGACCACCAT CACGACGGGT ACCAGGCACC GCCCCAAGAC   60
ATCGCGCTGC GGGTCAAGGC CTTGGAGTCT CTGCTGATCG AGAAAGGTCT TGTCGACCCA  120
GCGGCCATGG ACTTGTCGT CCAAACGTAT GAACACAAGG TAGGCCCCCG AAACGGCGCC   180
AAAGTCGTGG CCAAGGCCCTG GGTGGACCCT GCCTACAAGG CCCGTCTGCT GGCAGACGCA  240
ACTGCGGCAA TTGCCGAGCT GGGCTTCTCC GGGGTACAGG GCGAGGACAT GGTCAATTCTG  300
GAAAACACCC CCGCCGTCCA CAACGTCTTC GTTTGCACCT TGTGCTCTTG CTACCCATGG  360
CCGACGCTGG GCTTGCCCCC TGCCTGGTAC AAGGCCGCCG CCTACCGGTC CCGCATGGTG  420
AGCGACCCGC GTGGGGTTCT CGCGGAGTTC GGCCTGGTGA TCCCCGCCAA CAAGGAAATC  480
CGCGTCTGGG ACACCACGGC CGAATTGCGC TACATGGTGC TGCCGGAACG GCCCAGAACT  540
GAAGCCTACA GCGAAGAACA ACTGGCCGAA CTCGTTACCC GCGATTGAT GATCGGCACC  600
GGCCTGCCAA CCCAACCAC CCCATCTCAT TAA                               633

```

(2) INFORMATION FOR SEQ ID NO.:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 654 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:4:

```

ATGAATGGCA TTCACGATAC TGGCGGAGCA CATGGTTATG GGCCGGTTTA CAGAGAACCG   60
AACGAACCCG TCTTTCGCTA CGACTGGGAA AAAACGGTCA TGTCCCTGCT CCCGGCCCTG  120
CTCGCCAACG CGAACTTCAA CCTCGATGAA TTTCGGCATT CGATCGAGCG AATGGGCCCCG  180
GCCCCACTATC TGGAGGGGAA CTACTACGAA CACTGGCTTC ATGTCTTTGA GAACCTGCTG  240
GTCGAGAAGG GTGTGCTCAC GGCCACGGAA GTCGCGACCG GCAAGGCTGC GTCTGGCAAG  300
ACGGCGACGC GCGTGCTGAC GCCGGCCATC GTGGACGACT CGTCAGCACC GGGGCTTCTG  360
CGCCCGGGAG GAGGGTTCTC TTTTTCCT GTGGGGGACA AGGTCGCGT CCTCAACAAG  420
AACCCGGTGG GCCATACCCG CATGCCGCGC TACACGCGGG CAAAGTGGGG ACAGTGGTCA  480
TCGACCATGG TGTGTTTCGT GACGCCGGAC ACCGCGGCAC ACGGAAAGGG CGAGCAGCCC  540
CAGCACGTTT ACACCGTGAG TTTCACGTCG GTCGAACTGT GGGGGCAAGA CGCTTCCTCG  600
CCGAAGGACA CGATTCGCGT CGACTTGTGG GATGACTACC TGGAGCCAGC GTGA       654

```

(2) INFORMATION FOR SEQ ID NO.:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: unknown
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:5:

```
Gly Gln Ser His Thr His Asp His His His Asp Gly Tyr Gln Ala Pro
1          5          10          15
Pro Glu Asp Ile Ala Leu Arg Val Lys Ala Leu Glu Ser Leu
          20          25          30
```

(2) INFORMATION FOR SEQ ID NO.:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: unknown
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:6:

```
Asp Leu Val Val Gln Thr Tyr Glu His Lys Val Gly Pro
1          5          10
```

(2) INFORMATION FOR SEQ ID NO.:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: unknown
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:7:

```
Asn Gly Ala Lys Val Val Ala Lys Ala Trp Val Asp Pro Ala Tyr Lys
1          5          10          15
```

(2) INFORMATION FOR SEQ ID NO.:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: unknown
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:8:

Asp Pro Arg Gly Val Leu Ala Glu Phe Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO.:9:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: unknown
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:9:

Gly Leu Pro Thr Gln Pro Thr Pro Ser His
1 5 10

(2) INFORMATION FOR SEQ ID NO.:10:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 83 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: unknown
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:10:

Met Asn Gly Ile His Asp Thr Gly Gly Ala His Gly Tyr Gly Pro Val
1 5 10 15

Tyr Arg Glu Pro Asn Glu Pro Val Phe Arg Tyr Asp Trp Glu Lys Thr
20 25 30

Val Met Ser Leu Leu Pro Ala Leu Xaa Ala Asn Gly Asn Phe Asn Leu
35 40 45

Asp Glu Phe Arg His Ser Ile Glu Arg Met Gly Pro Ala His Tyr Leu
50 55 60

Glu Gly Thr Tyr Tyr Glu His Trp Leu His Val Phe Glu Asn Leu Leu
65 70 75 80

Val Glu Lys

(2) INFORMATION FOR SEQ ID NO.:11:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: unknown
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:11:

Gly Glu His Pro Gln His Val Tyr
1 5

(2) INFORMATION FOR SEQ ID NO.:12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: unknown
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:12:

Ser Phe Thr Ser Val Glu Leu Trp Gly Gln Asp Ala Ser Ser Pro Lys
1 5 10 15

(2) INFORMATION FOR SEQ ID NO.:13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: unknown
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:13:

Val Asp Leu Trp Asp Asp Tyr Leu Glu Pro Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO.:14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:14:

GGAATTCGAY CAYCAYCAYG A

21

(2) INFORMATION FOR SEQ ID NO.:15:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:15:

GGAATTCTTY TCCARTCRT A

21

(2) INFORMATION FOR SEQ ID NO.:16:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 726 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:16:

| | |
|--|-----|
| GAATTCGATC ACCATCACGA CGGGTACCAG GCACCGCCCG AAGACATCGC GCTGCGGGTC | 60 |
| AAGGCCTTGG AGTCTCTGCT GATCGAGAAA GGTCTTGTCTG ACCCAGCGGC CATGGACTTG | 120 |
| GTCGTCCAAA CGTATGAACA CAAGGTAGGC CCCCAGAAACG GCGCCAAAGT CGTGGCCAAG | 180 |
| GCCTGGGTGG ACCCTGCCTA CAAGGCCCGT CTGCTGGCAG ACGCAACTGC GGCAATTGCC | 240 |
| GAGCTGGGCT TCTCCGGGGT ACAGGGCGAG GACATGGTCA TTCTGGAAAA CACCCCCGCC | 300 |
| GTCCACAACG TCTTCGTTTG CACCTTGTGC TCTTGCTACC CATGGCCGAC GCTGGGCTTG | 360 |
| CCCCCTGCCT GGTACAAGGC CGCCGCCTAC CGGTCCCGCA TGGTGAGCGA CCCGCGTGGG | 420 |
| GTTCTCGCGG AGTTCGGCCT GGTGATCCCC GCCAACAAGG AAATCCGCGT CTGGGACACC | 480 |
| ACGGCCGAAT TGCCTACAT GGTGCTGCCG GAACGGCCCG GAACTGAAGC CTACAGCGAA | 540 |
| GAACAACTGG CCGAACTCGT TACCCGCGAT TCGATGATCG GCACCGGCCT GCCAACCCAA | 600 |
| CCCACCCCAT CTCATTAAGG AGTTCGTCAT GAATGGCATT CACGATACTG GCGGAGCACA | 660 |
| TGGTTATGGG CCGGTTTACA GAGAACCGAA CGAACCCGTC TTTCGCTACG ACTGGGAAAA | 720 |
| GAATTC | 726 |

(2) INFORMATION FOR SEQ ID NO.:17:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1440 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:17:

```

CGGGAGCGCA ATCTGCAAGG TGGCATTGGC CTTCACTGTC GATGCGGAGT TGAAGTCGCT      60
GTACCCCTTT TTTCAACCAC ACCAGGAGAA CCGCACCATG GGGCAATCAC ACACGCATGA      120
CCACCATCAC GACGGGTACC AGGCACCGCC CGAAGACATC GCGCTGCGGG TCAAGGCCTT      180
GGAGTCTCTG CTGATCGAGA AAGGTCTTGT CGACCCAGCG GCCATGGACT TGGTCGTCCA      240
AACGTATGAA CACAAGGTAG GCGCCGAAA CGGCGCCAAA GTCGTGGCCA AGGCCTGGGT      300
GGACCCTGCC TACAAGGCCC GTCTGCTGGC AGACGCAACT GCGGCAATTG CCGAGCTGGG      360
CTTCTCCGGG GTACAGGGCG AGGACATGGT CATTCTGGAA AACACCCCG CCGTCCACAA      420
CGTCTTCGTT TGCACCTTGT GCTCTTGCTA CCCATGGCCG ACGCTGGGCT TGCCCCCTGC      480
CTGGTACAAG GCCGCCGCCT ACCGGTCCCG CATGGTGAGC GACCCGCGTG GGGTTCTCGC      540
GGAGTTCGGC CTGGTGATCC CCGCCAACAA GGAAATCCGC GTCTGGGACA CCACGGCCGA      600
ATTGCGCTAC ATGGTGCTGC CGGAACGGCC CGGAAGTAA GCCTACAGCG AAGAACAAC      660
GGCCGAATC GTTACCCGCG ATTCGATGAT CGGCACCGGC CTGCCAACCC AACCCACCCC      720
ATCTCATTA GGAGTTCGTC ATGAATGGCA TTCACGATAC TGGCGGAGCA CATGGTTATG      780
GGCCGGTTTA CAGAGAACCG AACGAACCCG TCTTTCGCTA CGACTGGGAA AAAACGGTCA      840
TGTCCTTGCT CCCGGCCCTG CTCGCCAACG CGAACTTCAA CCTCGATGAA TTTCGGCATT      900
CGATCGAGCG AATGGGCCCCG GCCCACTATC TGGAGGGAAC CTACTACGAA CACTGGCTTC      960
ATGTCTTTGA GAACCTGCTG GTCGAGAAGG GTGTGCTCAC GGCCACGGAA GTCGCGACCG      1020
GCAAGGCTGC GTCTGGCAAG ACGGCGACGC GCGTGCTGAC GCCGGCCATC GTGGACGACT      1080
CGTCAGCACC GGGGCTTCTG CGCCCGGGAG GAGGGTTCTC TTTTTTTCCT GTGGGGGACA      1140
AGGTTCGCGT CCTCAACAAG AACCCGGTGG GCCATACCCG CATGCCGCGC TACACGCGGG      1200
CAAAGTGGGG ACAGTGGTCA TCGACCATGG TGTGTTTCGT GACGCCGGAC ACCGCGGCAC      1260
ACGGAAAGGG CGAGCAGCCC CAGCACGTTT ACACCGTGAG TTTCACGTCG GTCGAACTGT      1320
GGGGGCAAGA CGCTTCCTCG CCGAAGGACA CGATTGCGGT CGACTTGTGG GATGACTACC      1380
TGGAGCCAGC GTGATCATGA AAGACGAACG GTTTCATTG CCAGAGGGTT CGCTGAAGGA      1440

```

(2) INFORMATION FOR SEQ ID NO.:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:18:

GAGGAATTCA TGGCCATTAC TCGCCCTACC C 31

(2) INFORMATION FOR SEQ ID NO.:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:19:

GTCGAATTCT CAGAGCGTGC GCCAGTCCAC C 31

(2) INFORMATION FOR SEQ ID NO.:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1521 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: No

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:20:

ATGGCCATTA CTCGCCCTAC CCTCGACCAG GTTTTAGACA TCCGAACCCA GTTGACATG 60
 CAACTGACGC ACGAACAGGC AGCGTCCTAC CTGGAAGTGA TGCAACCGAG TTTCGACGCC 120
 TACGACCTGG TCGACGAACT GGCTGATTTT GTTCCGCCAA TACGCTACGA CCGCAGTTCA 180
 GGCTATCGCC ATCGGCCATC GGCCAAGGAA AACCCTCTGA ACGCCTGGTA CTACCGAACA 240
 GAAGTGAATG GTGCCCCGCA AGGCCTGCTG GCGGGCAAAA CCGTCGCGCT CAAAGATAAT 300
 ATCTCCCTGG CAGGCGTCCC CATGATGAAC GGCGCAGCGC CGTTGGAAGG CTTCGTCCCG 360
 GGGTTCGATG CCACGGTGGT CACCCGCTTG CTCGATGCGG GGGCGACCAT TCTCGGCAAA 420
 GCCACCTGCG AGCACTACTG CCTTTCAGGA GGCAGCCACA CCTCCGATCC AGCCCCGGTG 480
 CACAACCCAC ATCGCCACGG TTATGCCTCT GGCGGTTCTT CATCAGGCAG CGCGGCATTG 540
 GTTGCGTCCG GTGAGGTGGA CATCGCCGTG GGCGGCGATC AAGGCGGCTC CATTTCGGATC 600
 CCGTCGGCCT TCTGCGGTAC CTACGGCATG AAGCCCACCC ACGGCCTGGT GCCCTACACC 660
 GGCCTCATGG CGATTGAAGC CACGATCGAT CATGTCGGCC CCATCACCGG TAACGTGCGC 720
 GACAACGCGC TGATGCTGCA GGCAATGGCC GGTGCAGACG GACTCGACCC GCGCCAGGCG 780
 GCGCCTCAGG TCGATGACTA TTGCAGTTAC CTGGAAAAAG GCGTGAGCGG ACTCAGAATC 840
 GGGGTGTTGC AAGAGGGATT CGCGCTTGCT AACCAGGACC CTCGCGTGGC GGACAAAGTG 900

CGCGACGCCA TCGCCCGACT CGAGGCGTTG GCGGCTCATG TCGAGCCGGT CTCCATTCCC 960
 GAGCACAACC TGGCAGGGTT GTTGTGGCAC CCCATCGGTT GCGAAGGCTT GACCATGCAG 1020
 ATGATGCATG GCAACGGCGC AGGCTTTAAC TGGAAAGGAC TTTACGATGT CGGCCTGCTG 1080
 GACAAACAAG CCAGCTGGCG CGACGACGCA GACCAATTAT CCGCTCGCT CAAGCTCTGC 1140
 ATGTTCTGTCG GCCAATACGG CCTGTGCGCG TACAACGGAC GCTACTACGC CAAGGCCAG 1200
 AACCTTGAC GCTTTGCCCC GCAGGGATAC GACAAAGCGC TGCAAACCTA TGACCTGCTG 1260
 GTGATGCCGA CCACGCCCAT CACGGCCCCA CCCCACCCGC CAGCGAACTG CTCGATCAG 1320
 GAGTACGTGG CTCGCGCGTT GGAAATGATC GGCAATACCG CGCCACAGGA CATCACCGGG 1380
 CATCCGGCCA TGTCGATTCC GTGTGGCCTG CTGGACGGCC TGCCCSTCGG GCTGATGCTG 1440
 GTCGCAAAAC ACTACGCCGA GGGCACGATT TACCAAGCGG CGGCGGCGTT TGAAGCCTCG 1500
 GTGGACTGGC GCACGCTCTG A 1521

(2) INFORMATION FOR SEQ ID NO.:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 384 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
 - (B) STRAIN: P14K
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO.:21:

ATGGCCCTGT GTTTGACGAG CCTTGGCAGT CCCAGGCGTT TGCCTTGGTG GTCAGCATGC 60
 ACAAGGCCGG TCTCTTTCAG TGGAAAGACT GGGCCGAGAC CTTACCCGCC GAAATCGACG 120
 CTTCCCCGCT CTGCCGGCGA AAGCGTCAAC GACACCTACT ACCGGCAATG GGTGTCGGCG 180
 CTGGAAAAGT TGGTGGCGTC GCTGGGGCTT GTGACGGGTG GAGACGTCAA CTCGCGCGCA 240
 CAGGAGTGGA AACAGGCCCA CCTCAACACC CCACATGGGC ACCCGATCCT GCTGGCCCAT 300
 GCGCTTTGCC CGCCAGCGAT CGACCCCAAG CACAAGCACG AGCCACAACG CTCACCGATC 360
 AAGGTCGTTG CCGCAATGGC TTGA 384

(2) INFORMATION FOR SEQ ID NO.:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 127 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown
- (ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
(B) STRAIN: P14K

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:22:

```

Met Ala Leu Cys Leu Thr Ser Leu Gly Ser Pro Arg Arg Leu Pro Trp
1           5           10           15
Trp Ser Ala Cys Thr Arg Pro Val Ser Phe Ser Gly Lys Thr Gly Pro
20           25           30
Arg Pro Ser Pro Pro Lys Ser Thr Leu Pro Arg Ser Ala Gly Glu Ser
35           40           45
Val Asn Asp Thr Tyr Tyr Arg Gln Trp Val Ser Ala Leu Glu Lys Leu
50           55           60
Val Ala Ser Leu Gly Leu Val Thr Gly Gly Asp Val Asn Ser Arg Ala
65           70           75           80
Gln Glu Trp Lys Gln Ala His Leu Asn Thr Pro His Gly His Pro Ile
85           90           95
Leu Leu Ala His Ala Leu Cys Pro Pro Ala Ile Asp Pro Lys His Lys
100          105          110
His Glu Pro Gln Arg Ser Pro Ile Lys Val Val Ala Ala Met Ala
115          120          125

```

(2) INFORMATION FOR SEQ ID NO.:23:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:23:

GATGCGGCCA TAGGCGAATT C

21

(2) INFORMATION FOR SEQ ID NO.:24:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:24:

ACCGCCACCG ACTACCTGCA G

21

(2) INFORMATION FOR SEQ ID NO.:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:25:

GTCAGCCTGA GCAATCTGCA G

21

(2) INFORMATION FOR SEQ ID NO.:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:26:

GAATTCGGAA AAAATCGTAC G

21

(2) INFORMATION FOR SEQ ID NO.:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1401 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (B) STRAIN: AMIDASE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:27:

| | |
|---|-----|
| ATGAGTTCGC TAACCCGCCT CACCCTCGCG CAAGTTGCGC AGAAACTTAA GGCACGGGAA | 60 |
| GTCTCCGCCG TTGAAGTTCT GGACGCCTGT CTGACGCAGG TGCCTCCAC CGAAAAACAG | 120 |
| ATCAGTGCCT ACGTGTGCGT GCTGGAGGAT CAGGCCCGTG CAGCAGCCCA CGCAACTGAC | 180 |
| GCCGACATCC GCGGGCGCTG GAAAGGCCCG CTGCATGGCG TGCCTGTAGC GGTCAAGGAC | 240 |
| TTATACGACA TCGCTGGCGT ACCCACCACG GCATCGTCGC CAGCGCACGA ATTGGACGCG | 300 |
| CAGCAAGACC CGGCTAGAGT CCGGCGCTTA CAAGACGCAG GTGCCGTTAT CCTTGCAAG | 360 |
| ACCCATACGC ACGAATTCGC CTATGGCCGC ATCACTCCGA AGTCGCGCAA CCCCAGGGAC | 420 |
| CCGGGAAGAA CACCGGGTGG CTCCAGCGGC GGCTCGGCGG CCACGGTCGC AGCCTGCTGC | 480 |
| GTCTACTTGG CGACCGGCAC CGACACCGGT GGATCCGTTT GCAICCTTC GTCGATGTGC | 540 |

```

AACACCGTAG GCCTGAAGCA ACCTACGGTC GGCCGCGTGC ACGGTGCCGG TGTGAGTTCA 600
CTTTCCTGGA GCCTGGACCA TCCAGGCCCG ATCACGCGCA CCGTGGAAGA CACGGCGCTC 660
ATGCTTCAGG TGATGGCTGG CTTCGATCCA GCCGACCCGC GGTCGTTGGA TGAGCCGGTG 720
CCCAGCTATG CCGAAGGGCT CGGCCAAGGC GTGAAAGGCC TGCCTGGGG TGTGCCGAAG 780
AACTACTTCT TCGACCGCGT GGACCCGGAA GTTGAAAGTG CGGTTCGTGC CGCCATCGAT 840
CAACTGAAAG AGCTGGGCGC CGAACTGGTG GAAGTCGAAG TGCCCATGGC CGAGCAGATC 900
ATCCCGGTGA AGTTCGGGAT CATGCTACCC GAAGCCAGCG CCTACCACCG CACGATGCTG 960
CGCGAGTCAC CCGAGCTCTA CACCGCCGAT GTCCGCATAC TGCTGGAAC TCGGAGATCTA 1020
GTCACCGCCA CCGACTACCT GCAGGCGCAG CGCGTCCGTA CGCTGATGCA GCGCGCGGTG 1080
GCCGAGATGT ACCAGCGCAT CGATGTGCTG ATCGCACCCA CACTGCCCAT CCCGGCTGCT 1140
CGCAGCGGGG AGGAGGTCCA CACATGGCCG GACGGCACGG TAGAGGCGTT GGTCATGGCC 1200
TATACGCGCT TCACCTCGTT CGGCAACGTG ACAGGATTAC CCACGCTGAA CCTGCCCTGT 1260
GGTTTCTCCA AGGATGGGTT GCGATCGGCA TGCAGATCAG GCCGGCCGCT GGACGAGAAG 1320
ACCCTGCTGC GTGCTGGGCT GGCCTACGAG AAAGCCACGA CCTGGCACCA GCGTCATCCG 1380
GAACTGATCG GAGCGGGCTG A 1401

```

(2) INFORMATION FOR SEQ ID NO.:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 466 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (B) STRAIN: AMIDASE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.:28:

```

Met Ser Ser Leu Thr Arg Leu Thr Leu Ala Gln Val Ala Gln Lys Leu
1           5           10           15
Lys Ala Arg Glu Val Ser Ala Val Glu Val Leu Asp Ala Cys Leu Thr
20           25           30
Gln Val Arg Ser Thr Glu Lys Gln Ile Ser Ala Tyr Val Cys Val Leu
35           40           45
Glu Asp Gln Ala Arg Ala Ala Ala His Ala Thr Asp Ala Asp Ile Arg
50           55           60
Gly Arg Trp Lys Gly Pro Leu His Gly Val Pro Val Ala Val Lys Asp
65           70           75           80

```

Leu Tyr Asp Ile Ala Gly Val Pro Thr Thr Ala Ser Ser Pro Ala His
 85 90 95
 Glu Leu Asp Ala Gln Gln Asp Pro Ala Arg Val Arg Arg Leu Gln Asp
 100 105 110
 Ala Gly Ala Val Ile Leu Gly Lys Thr His Thr His Glu Phe Ala Tyr
 115 120 125
 Gly Arg Ile Thr Pro Lys Ser Arg Asn Pro Arg Asp Pro Gly Arg Thr
 130 135 140
 Pro Gly Gly Ser Ser Gly Gly Ser Ala Ala Thr Val Ala Ala Cys Cys
 145 150 155 160
 Val Tyr Leu Ala Thr Gly Thr Asp Thr Gly Gly Ser Val Arg Ile Pro
 165 170 175
 Ser Ser Met Cys Asn Thr Val Gly Leu Lys Gln Pro Thr Val Gly Arg
 180 185 190
 Val His Gly Ala Gly Val Ser Ser Leu Ser Trp Ser Leu Asp His Pro
 195 200 205
 Gly Pro Ile Thr Arg Thr Val Glu Asp Thr Ala Leu Met Leu Gln Val
 210 215 220
 Met Ala Gly Phe Asp Pro Ala Asp Pro Arg Ser Leu Asp Glu Pro Val
 225 230 235 240
 Pro Ser Tyr Ala Glu Gly Leu Gly Gln Gly Val Lys Gly Leu Arg Trp
 245 250 255
 Gly Val Pro Lys Asn Tyr Phe Phe Asp Arg Val Asp Pro Glu Val Glu
 260 265 270
 Ser Ala Val Arg Ala Ala Ile Asp Gln Leu Lys Glu Leu Gly Ala Glu
 275 280 285
 Leu Val Glu Val Glu Val Pro Met Ala Glu Gln Ile Ile Pro Val Lys
 290 295 300
 Phe Gly Ile Met Leu Pro Glu Ala Ser Ala Tyr His Arg Thr Met Leu
 305 310 315 320
 Arg Glu Ser Pro Glu Leu Tyr Thr Ala Asp Val Arg Ile Leu Leu Glu
 325 330 335
 Leu Gly Asp Leu Val Thr Ala Thr Asp Tyr Leu Gln Ala Gln Arg Val
 340 345 350
 Arg Thr Leu Met Gln Arg Ala Val Ala Glu Met Tyr Gln Arg Ile Asp
 355 360 365
 Val Leu Ile Ala Pro Thr Leu Pro Ile Pro Ala Ala Arg Ser Gly Glu
 370 375 380
 Glu Val His Thr Trp Pro Asp Gly Thr Val Glu Ala Leu Val Met Ala
 385 390 395 400
 Tyr Thr Arg Phe Thr Ser Phe Gly Asn Val Thr Gly Leu Pro Thr Leu
 405 410 415

Asn Leu Pro Cys Gly Phe Ser Lys Asp Gly Leu Arg Ser Ala Cys Arg
420 425 430

Ser Gly Arg Pro Leu Asp Glu Lys Thr Leu Leu Arg Ala Gly Leu Ala
435 440 445

Tyr Glu Lys Ala Thr Thr Trp His Gln Arg His Pro Glu Leu Ile Gly
450 455 460

Ala Gly
465

WHAT IS CLAIMED IS:

1. A nucleic acid fragment encoding the α subunit of a stereospecific nitrile hydratase enzyme, said nucleic acid fragment having at least a 64% base
5 homology with the α subunit coding region of the *Rhodococcus rhodochrous* J1 L-NHase gene and said enzyme capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.
2. A nucleic acid fragment encoding the β
10 subunit of a stereospecific nitrile hydratase enzyme, said nucleic acid fragment having at least a 52% base homology with the β subunit coding region of the *Rhodococcus rhodochrous* J1 L-NHase gene and said enzyme capable of catalyzing the hydrolysis of racemic aryl-2-
15 alkane nitriles to the corresponding R- or S- amides.
3. A nucleic acid fragment comprising the nucleic acid fragments of Claims 1 and 2 encoding a stereospecific nitrile hydratase enzyme, said enzyme capable of catalyzing the hydrolysis of racemic aryl-2-
20 alkane nitriles to the corresponding R- or S- amides.
4. A nucleic acid fragment encoding the α subunit of a stereospecific nitrile hydratase enzyme, said nucleic acid fragment having the nucleotide sequence as represented in SEQ ID NO.:3 and said enzyme
25 capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.
5. A nucleic acid fragment encoding the β subunit of a stereospecific nitrile hydratase enzyme, said nucleic acid fragment having the nucleotide
30 sequence as represented in SEQ ID NO.:4 and said enzyme capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.
6. A nucleic acid fragment encoding both the α and β subunits of a stereospecific nitrile hydratase
35 enzyme, said nucleic acid fragment having the nucleotide sequence as represented in SEQ ID NO.:17 and said enzyme capable of catalyzing the hydrolysis of

racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.

7. A polypeptide α subunit of a stereospecific nitrile hydratase enzyme, said α subunit having the amino acid sequence as represented in SEQ ID NO.:1 and said enzyme catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.

8. A polypeptide β subunit of a stereospecific nitrile hydratase enzyme, said β subunit having the amino acid sequence as represented in SEQ ID NO.:2 and said enzyme catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.

9. A stereospecific nitrile hydratase enzyme comprising 1) the polypeptide α subunit of Claim 7 and 2) the polypeptide β subunit of Claim 8 in proper conformation such that said enzyme catalyzes the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.

10. A 6.5 kb nucleic acid fragment encoding a nitrile hydratase enzyme and the accessory nucleic acid fragments necessary for the enzymes's active expression as characterized by the restriction map shown in Figure 2.

11. A recombinant expression vector comprising the 6.5 kb nucleic acid fragment of Claim 10, said vector capable of transforming a suitable host cell for the expression of active stereospecific nitrile hydratase as characterized by the restriction map shown in Figure 3.

12. A transformed microbial host cell comprising the nucleic acid fragment of Claim 5 wherein said host cell expresses active nitrile hydratase enzyme capable of catalyzing the hydrolysis of racemic aryl-2 nitriles to the corresponding R- or S- amides.

13. A transformed microbial host cell comprising the 6.5 kb nucleic acid fragment of Claim 10 wherein

said host cell expresses active nitrile hydratase enzyme capable of catalyzing the hydrolysis of racemic aryl-2 nitriles to the corresponding R- or S- amides.

14. The transformed host cell of Claims 12 or 13 wherein said host cell is selected from the group consisting of bacteria, yeast, and filamentous fungi.

15. The transformed host cell of Claim 14 wherein said host cell is a bacterium selected from the group consisting of the genera *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, and *Streptomyces*.

16. A transformed host cell of Claim 14 wherein said host cell is a yeast selected from the group consisting of the genera *Pichia*, *Hansenula*, and *Saccharomyces*.

17. A transformed host cell of Claim 14 wherein said host cell is a filamentous fungus selected from the group consisting of the genera *Aspergillus*, *Neurospora*, and *Penicillium*.


18. A transformed host cell of Claim 15 wherein said host cell is *Escherichia coli*.

19. An expression vector comprising 1) a 5.0 kb nucleic acid fragment of the 6.5 kb nucleic acid fragment of Claim 10, and 2) a nucleic acid fragment having the nucleic acid sequence as given in SEQ ID NO.:20, wherein said nucleic acid fragment encodes an amidase enzyme, and wherein said expression vector is capable of transforming suitable host cells for the co-expression of active stereospecific nitrile hydratase and amidase.

20. A transformed host cell comprising the vector of Claim 19.





21. A transformed host cell of Claim 20 wherein said host is selected from the group consisting of the genera *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Streptomyces*, *Hansenula*, *Saccharomyces*, *Pichia*, *Aspergillus*, *Neurospora*, and *Penicillium*.


22. A method for converting a nitrile of the formula



wherein:

A is selected from the group consisting of:






A-10

A-11

R^1 is C_1-C_4 alkyl;

R^2 is H or OH;


R^3 is H, Cl, OCF_2H , $(CH_3)_2CHCH_2$, $H_2C=C(CH_3)CH_2NH$,



R^4 is Cl or F;





- 5 to the corresponding amide comprising
 - 1) contacting said nitrile with the transformed host cell comprising a nucleic acid fragment having the nucleotide sequence represented by SEQ ID NO.:17 that stereospecifically converts the
 - 10 racemic nitrile to the corresponding enantiomeric R- or S-amide, the host cell selected from the group consisting of *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Streptomyces*, *Hansenula*, *Saccharomyces*, *Pichia*, *Aspergillus*, *Neurospora*, and
 - 15 *Penicillium*,
 - 2) recovering the product of Step 1).


23. A method for converting a nitrile of the formula



wherein:

A is selected from the group consisting of:






A-10

A-11

R^1 is C_1 - C_4 alkyl;

R^2 is H or OH;

R^3 is H, Cl, OCF_2H , $(CH_3)_2CHCH_2$, $H_2C=C(CH_3)CH_2NH$,



R^4 is Cl or F;

- 5 to the corresponding enantiomeric (R) or (S)-carboxylic acid comprising

- 1) contacting the nitrile with the transformed host of Claim 20 that stereospecifically converts the racemic nitrile to the corresponding enantiomeric (R)- or (S)-carboxylic acid, the host cell selected from the group consisting of *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Streptomyces*, *Hansenula*, *Saccharomyces*, *Pichia*, *Aspergillus*, *Neurospora*, and *Penicillium*, and
- 15 2) recovering the product of Step 1).

24. A nucleic acid fragment encoding the α and β subunits of a stereospecific nitrile hydratase enzyme, said portion of the nucleic acid fragment encoding the α subunit having at least a 64% base homology to the *Rhodochrous* J1 L-NHase gene and said portion of the nucleic acid fragment encoding the β subunit having at least a 52% base homology to the *Rhodochrous* J1 L-NHase gene, said enzyme catalyzing the hydrolysis of racemic

aryl-2-alkane nitriles to the corresponding R- or S-amides.

25. The polypeptide encoded by any one of the nucleic acid fragments of Claims 1, 2, 6, or 24.

5 26. A transformed *Escherichia coli* comprising pSW2 and identified as ATCC 69888.

27. A transformed *Escherichia coli* comprising pSW17 and identified as ATCC 69889.

10 28. A transformed host cell comprising the nucleic acid fragments of Claims 1 and 2, wherein said transformed host expresses an active nitrile hydratase.

29. A nucleic acid fragment corresponding to the P14K region of the 6.5 kb fragment as shown in Figure 10 wherein said fragment encodes a polypeptide
15 having an amino acid sequence as represented in SEQ ID NO.:22.

30. A nucleic acid fragment of Claim 29 where said fragment has the base sequence as represented in SEQ ID NO.:21.

20 32. A P14K polypeptide encoded by the nucleic acid fragment of Claim 29 and having the amino acid sequence as represented in SEQ ID NO.:22, wherein the expression of said polypeptide is required for the bioactivity of expressed *Pseudomonas putida* NRRL-18668
25 stereospecific nitrile hydratase capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S- amides.

33. A nucleic acid fragment encoding a 18668 amidase having an amino acid sequence as represented in
30 SEQ ID NO.:28, wherein said amino acid sequence may encompass amino acid substitutions, deletions or additions that do not alter the function of said amidase.

34. A nucleic acid fragment of Claim 33 where
35 said fragment has the base sequence as given by SEQ ID NO.:27.

35. An 18668 amidase polypeptide encoded by the nucleic acid fragment of Claim 33 and having the amino

acid sequence as represented in SEQ ID NO.:28, wherein said amino acid sequence may encompass amino acid substitutions, deletions or additions that do not alter the function of said amidase.

5 36. A transformed microbial host containing the nucleic acid fragments of Claim 6 and Claim 29 said host capable of catalyzing the hydrolysis of racemic aryl-2-alkane nitriles to the corresponding R- or S-amides.

10 37. A transformed host according to Claim 36 wherein said host is a member of the genus *Pichia*.


 38. A transformed microbial host containing the nucleic acid fragment of Claim 6, Claim 29 and Claim 33 said host capable of catalyzing the hydrolysis of
15 racemic aryl-2-alkane nitriles to the corresponding R- or S- amides and the conversion of said amides to the corresponding enantiomeric (R) or (S)-carboxylic acids.

 39. A transformed *Pichia pastoris* comprising pSW50 and identified as ATCC 74391.

20 41. A transformed *E. coli* comprising pSW37 and identified as ATCC 98174.


 42. A transformed *E. coli* comprising pSW23 and identified as ATCC 98175.

25 43. A method for converting a nitrile of the formula



wherein:


A is selected from the group consisting of:




A-1

A-2


A-3




A-4




A-5




A-6




A-7



A-8




A-9



A-10

and





A-11

R^1 is C_1 - C_4 alkyl;

R^2 is H or OH;

R^3 is H, Cl, OCF_2H , $(CH_3)_2CHCH_2$, $H_2C=C(CH_3)CH_2NH$,






R^4 is Cl or F;

to the corresponding amide comprising

- 1) contacting said nitrile with a transformed host cell of Claim 36 comprising the nucleic acid fragments of Claim 6 and Claim 29, the host cell capable of stereospecifically converting the racemic nitrile to the corresponding enantiomeric R- or S-amide, the host cell selected from the group consisting of *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Streptomyces*, *Hansenula*, *Saccharomyces*, *Pichia*, *Aspergillus*, *Neurospora*, and *Penicillium*,


- 2) recovering the product of Step 1).


44. A method for converting a nitrile of the formula




wherein:

A is selected from the group consisting of:







A-4




A-5




A-6




A-7




A-8



A-9



A-10








A-11


R¹ is C₁-C₄ alkyl;

R² is H or OH;

R³ is H, Cl, OCF₂H, (CH₃)₂CHCH₂, H₂C=C(CH₃)CH₂NH,



or



; and

R⁴ is Cl or F;

5 to the corresponding enantiomeric (R) or (S)-carboxylic acid comprising

- 1) contacting the nitrile with the transformed host cell of Claim 39 that stereo-specifically converts the racemic nitrile to the corresponding enantiomeric (R)- or (S)-carboxylic acid,
- 5 the host cell selected from the group consisting of *Escherichia*, *Pseudomonas*, *Rhodococcus*, *Acinetobacter*, *Bacillus*, *Streptomyces*, *Hansenula*, *Saccharomyces*, *Pichia*, *Aspergillus*, *Neurospora*, and *Penicillium*, and
- 2) recovering the product of Step 1).

10

1/10

FIG. 1





FIG. 2

FIG. 3





FIG. 4







FIG.5

4/10

FIG. 6



5/10
FIG. 7

SEQ ID NO:17:

| | |
|--|------|
| CGGGAGCGCA ATCTGCAAGG TGGCATTGGC CTTCACTGTC GATGCCGAGT TGAAGTCGCT | 60 |
| GTACCCCTTT TTTCAACCAC ACCAGGAGAA CCGCACCATG GGGCAATCAC ACACGCATGA | 120 |
| CCACCATCAC GACGGGTACC AGGCACCGCC CGAAGACATC GCGCTGCGGG TCAAGGCCTT | 180 |
| GGAGTCTCTG CTGATCGAGA AAGGTCTTGT CGACCCAGCG GCCATGGACT TGGTCGTCCA | 240 |
| AACGTATGAA CACAAGGTAG GCCCCGAAA CGGCGCCAAA GTCGTGGCCA AGGCCTGGGT | 300 |
| GGACCCTGCC TACAAGGCC GTCTGCTGGC AGACGCAACT GCGGCAATTG CCGAGCTGGG | 360 |
| CTTCTCCGGG GTACAGGGCG AGGACATGGT CATTCTGGAA AACACCCCCG CCGTCCACAA | 420 |
| CGTCTTCGTT TGCACCTTGT GCTCTTGCTA CCCATGGCCG ACGCTGGGCT TGCCCCCTGC | 480 |
| CTGGTACAAG GCCGCCGCCT ACCGGTCCCG CATGGTGAGC GACCCGCGTG GGGTTCTCGC | 540 |
| GGAGTTCGGC CTGGTGATCC CCGCCAACAA GGAAATCCGC GTCTGGGACA CCACGGCCGA | 600 |
| ATTGCGCTAC ATGGTGCTGC CGGAACGGCC CGGAACTGAA GCCTACAGCG AAGAACAAC | 660 |
| GGCCGAACCTC GTTACCCGCG ATTCGATGAT CGGCACCGGC CTGCCAACCC AACCACCCC | 720 |
| ATCTCATTA AAGAGTTCGTC ATGAATGGCA TTCACGATAC TGGCGGAGCA CATGGTTATG | 780 |
| GGCCGGTTTA CAGAGAACCG AACGAACCCG TCTTTGCTA CGACTGGGAA AAAACGGTCA | 840 |
| TGTCCCTGCT CCCGGCCCTG CTCGCCAACG CGAACTTCAA CCTCGATGAA TTTCGGCATT | 900 |
| CGATCGAGCG AATGGGCCCC GCCCACTATC TGGAGGGAAC CTACTACGAA CACTGGCTTC | 960 |
| ATGTCTTTGA GAACCTGCTG GTCGAGAAGG GTGTGCTCAC GGCCACGGAA GTCGCGACCG | 1020 |
| GCAAGGCTGC GTCTGGCAAG ACGGCGACGC GCGTGCTGAC GCCGGCCATC GTGGACGACT | 1080 |
| CGTCAGCACC GGGGCTTCTG CGCCCGGGAG GAGGGTTCTC TTTTTTCTCT GTGGGGGACA | 1140 |
| AGGTTTCGCGT CCTCAACAAG AACCCTGGTG GCCATACCCG CATGCCGCGC TACACGCGGG | 1200 |
| CAAAGTGGGG ACAGTGGTCA TCGACCATGG TGTGTTTCGT GACGCCGGAC ACCGCGGCAC | 1260 |
| ACGGAAAGGG CGAGCAGCCC CAGCACGTTT ACACCGTGAG TTTCACGTCG GTCGAACTGT | 1320 |
| GGGGGCAAGA CGCTTCCTCG CCGAAGGACA CGATTCGCGT CGACTTGTGG GATGACTACC | 1380 |
| TGGAGCCAGC GTGATCATGA AAGACGAACG GTTTCATTG CCAGAGGGTT CGCTGAAGGA | 1440 |

6/10




FIG. 8




FIG. 9




FIG. 10

7/10

FIG. 11





FIG. 12



8/10

FIG. 13





FIG. 14



9/10

FIG. 15





FIG. 16



10/10

FIG. 17





FIG. 18



SUBSTITUTE SHEET (RULE 26)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|---|
| <p>(51) International Patent Classification ⁶ : C12N 9/78, 15/55, 15/31, C07K 14/21, C12N 15/63, 15/62, 1/19, 1/21, C12P 13/02, 7/40 // (C12N 9/78, C12R 1:40) (C12N 1/19, C12R 1:84)</p> | <p>A3</p> | <p>(11) International Publication Number: WO 97/12964 (43) International Publication Date: 10 April 1997 (10.04.97)</p> |
| <p>(21) International Application Number: PCT/US96/15969 (22) International Filing Date: 3 October 1996 (03.10.96) (30) Priority Data: 60/004,914 6 October 1995 (06.10.95) US (71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FALLON, Robert, Donald [US/US]; 104 Black Oak Drive, Elkton, MD 21921- 2022 (US). NELSON, Mark, James [US/US]; 707 Fiske Lane, Newark, DE 19711-3133 (US). PAYNE, Mark, Scott [US/US]; 2408 Newell Drive, Wilmington, DE 19808-3344 (US). (74) Agents: FLOYD, Linda, A. et al.; E.I. du Pont de Nemours and Company, Legal Patent Record Center, 1007 Market Street, Wilmington, DE 19898 (US).</p> | | <p>(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 19 June 1997 (19.06.97)</p> |
| <p>(54) Title: NUCLEIC ACID FRAGMENTS ENCODING STEREOSPECIFIC NITRILE HYDRATASE AND AMIDASE ENZYMES AND RECOMBINANT MICROORGANISMS EXPRESSING THOSE ENZYMES USEFUL FOR THE PRODUCTION OF CHIRAL AMIDES AND ACIDES</p> | | |
| | | |
| <p>(57) Abstract</p> <p>The present invention provides a nitrile hydratase nucleic acid fragment isolated from <i>Pseudomonas putida</i> which encodes a nitrile hydratase activity capable of catalyzing the hydrolysis of certain racemic nitriles to the corresponding R- or S- amides. Also provided are transformed microorganisms capable of the active expression of said nitrile hydratase activity. Additionally, the invention provides a transformant harboring the nitrile hydratase gene in conjunction with an amidase gene, both of which may be co-expressed producing active nitrile hydratase and amidase enzymes respectively. Methods for the production of such enantiomeric materials are also provided.</p> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|---------------------------------------|----|--------------------------|
| AM | Armenia | GB | United Kingdom | MW | Malawi |
| AT | Austria | GE | Georgia | MX | Mexico |
| AU | Australia | GN | Guinea | NE | Niger |
| BB | Barbados | GR | Greece | NL | Netherlands |
| BE | Belgium | HU | Hungary | NO | Norway |
| BF | Burkina Faso | IE | Ireland | NZ | New Zealand |
| BG | Bulgaria | IT | Italy | PL | Poland |
| BJ | Benin | JP | Japan | PT | Portugal |
| BR | Brazil | KE | Kenya | RO | Romania |
| BY | Belarus | KG | Kyrgyzstan | RU | Russian Federation |
| CA | Canada | KP | Democratic People's Republic of Korea | SD | Sudan |
| CF | Central African Republic | KR | Republic of Korea | SE | Sweden |
| CG | Congo | KZ | Kazakhstan | SG | Singapore |
| CH | Switzerland | LI | Liechtenstein | SI | Slovenia |
| CI | Côte d'Ivoire | LK | Sri Lanka | SK | Slovakia |
| CM | Cameroon | LR | Liberia | SN | Senegal |
| CN | China | LT | Lithuania | SZ | Swaziland |
| CS | Czechoslovakia | LU | Luxembourg | TD | Chad |
| CZ | Czech Republic | LV | Latvia | TG | Togo |
| DE | Germany | MC | Monaco | TJ | Tajikistan |
| DK | Denmark | MD | Republic of Moldova | TT | Trinidad and Tobago |
| EE | Estonia | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | UG | Uganda |
| FI | Finland | MN | Mongolia | US | United States of America |
| FR | France | MR | Mauritania | UZ | Uzbekistan |
| GA | Gabon | | | VN | Viet Nam |

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 96/15969

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N9/78 C12N15/55 C12N15/31 C07K14/21 C12N15/63
C12N15/62 C12N1/19 C12N1/21 C12P13/02 C12P7/40
//(C12N9/78,C12R1:40),(C12N1/19,C12R1:84)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C12P C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|--------------------------|
| A | WO 92 05275 A (DU PONT DE NEMOURS AND COMPANY) 2 April 1992 cited in the application see Examples 1-4 --- | 1-39, 41-44 |
| A | JP 06 303 971 A (MITSUBISHI KASEI CORP.) 1 November 1994 see SEQ ID Nos 1 and 2 --- | 1-28, 36-39, 42-44 |
| A | EP 0 445 646 A (NITTO CHEMICAL INDUSTRY CO., LTD) 11 September 1991 see SEQ ID Nos 3 and 4 --- | 1-28, 36-39, 42-44 |
| | -/-- | |



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

25 April 1997

Date of mailing of the international search report

16. 05. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Alt, G

INTERNATIONAL SEARCH REPORT

Int. Patent Application No.
PCT/US 96/15969

| C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT | | |
|--|--|--|
| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| A | EP 0 579 907 A (MITSUBISHI KASEI CORP.) 26 January 1994 see SEQ ID No. 1 --- | 1,3,4,6, 7,9-28, 36-39, 42-44 |
| A | BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 1129, 1991, pages 23-33, XP000653047 KOBAYASHI, M. ET AL.: "Cloning, nucleotide sequence and expression in Escherichia coli of two cobalt-containing nitrile hydratase genes from Rhodococcus rhodochrous J1" cited in the application see the whole document --- | 1-32, 36-39, 41-44 |
| A | JOURNAL OF BACTERIOLOGY, vol. 173, no. 8, April 1991, pages 2465-2472, XP000653038 NISHIYAMA, M. ET AL.: "Cloning and characterization of genes responsible for metabolism of nitrile compounds from Pseudomonas chlororaphis B23" cited in the application see Figure 2; page 2470 --- | 19,29-32 |
| A | BIOSCIENCE, BIOTECHNOLOGY, AND BIOCHEMISTRY , vol. 58, October 1994, pages 1859-1865, XP000653085 HASHIMOTO, Y. ET AL.: "Nitrile hydratase gene form Rhodococcus sp. N-774 requirement for its downstream region for efficient expression" see the whole document --- | 29-32 |
| A | EP 0 433 117 A (RHONE-POULENC SANTE) 19 June 1991 see Figure 8 --- | 33-35 |
| A | JOURNAL OF BACTERIOLOGY, vol. 172, no. 12, 1990, pages 6764-6773, XP000318450 MAYAUX, J.-F. ET AL.: "Purification , cloning and primary structure of an enantiomer-selective amidase from Brevibacterium sp. Strain R312: Structural Evidence fro genetic coupling with nitrile hydratase" see Figure 2 ----- | 33-35 |

Form PCT ISA 210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. l. Application No.

PCT/US 96/15969

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|---|--|
| WO 9205275 A | 02-04-92 | AT 134706 T DE 69117521 D DE 69117521 T EP 0549710 A EP 0683145 A EP 0683235 A ES 2085487 T HK 122196 A JP 6506343 T US 5593871 A | 15-03-96 04-04-96 19-09-96 07-07-93 22-11-95 22-11-95 01-06-96 19-07-96 21-07-94 14-01-97 |
| JP 6303971 A | 01-11-94 | NONE | |
| EP 445646 A | 11-09-91 | AT 138101 T AU 627648 B AU 7129391 A CA 2037291 A CN 1054616 A DE 69119454 D DE 69119454 T ES 2089044 T JP 4211379 A KR 9502004 B PL 166201 B | 15-06-96 27-08-92 12-09-91 29-08-91 18-09-91 20-06-96 26-09-96 01-10-96 03-08-92 08-03-95 28-04-95 |
| EP 579907 A | 26-01-94 | AU 3692593 A CN 1084565 A JP 6025296 A US 5314819 A | 04-11-93 30-03-94 01-02-94 24-05-94 |
| EP 433117 A | 19-06-91 | FR 2655660 A AU 631696 B AU 6661490 A CA 2030073 A CN 1052508 A JP 4218379 A US 5260208 A | 14-06-91 03-12-92 13-06-91 12-06-91 26-06-91 07-08-92 09-11-93 |